

LAKE COUNTY, IL

2016 INDEPENDENCE GROVE SUMMARY REPORT

LAKE COUNTY HEALTH DEPARTMENT
ECOLOGICAL SERVICES



Independence Grove

Independence Grove is a borrow pit lake that was purchased in 1978 by the Lake County Forest Preserve. After mining the quarry of remaining resources, establishing a safe shoreline and wildlife habitat, the lake was open for public use in spring of 2001. The lake has a surface area of approximately 129.3 acres, a maximum depth of 52.0 feet, and an estimated average depth of 15 feet. It offers many amenities such as boating (rentals only), fishing, picnic areas, hiking trails, swimming beach, and aesthetics. Electric motors are allowed on the lake.

Water levels were taken from a foot bridge at the east entrance to North Bay. The water level decreased approximately 12.72 inches throughout the season (May to September). Independence Grove was stratified for most of the summer with a thermocline forming between 14 to 20 feet. Dissolved oxygen concentrations were adequate to support aquatic life (> 5 mg/L) in the epilimnion for the entire summer, however the hypolimnion had anoxic conditions (<1 mg/L) June through September.

The water quality of Independence Grove was very good. Total suspended solid concentrations (TSS) fluctuated throughout the season with a maximum concentration of 2.2 mg/L in August and September. The average TSS concentration for the season was 1.6

ECOLOGICAL SERVICES WATER QUALITY SPECIALISTS

Gerard Urbanozo

ECOLOGICAL SERVICES

Alana Bartolai

gurbanozo@lakecountyil.gov

Phone: 847-377-8030

Abartolai2@lakecountyil.gov

INSIDE THIS ISSUE:

Summary	1
Watershed	2
Watershed	3
Land Use	
VLMP	4
Water Quality	5
Total Suspended Solids	
Nutrients	6
Conductivity - Chloride	
Trophic State Index	7
Lake Level	
Bathymetric Maps	8
Stratification	
Blue-Green Algae	9
FQI	10
Aquatic Plants	11
Common Aquatic Plants	12
Common Aquatic Plants	13
EWM	14
Curlyleaf Pondweed	15
Aquatic Herbicide	16
Coarse Woody Habitat	
Manual Removal of Plants	17
Protect Your Waters	
Dissolved Oxygen	18
Aquatic Plants and Fish	19
Shoreline Erosion	20
Zebra Mussels	21
Swimming Beach	22
Lake Management Plans	23
Lake Recommendations	24

See Figures and Tables
 In Appendix A and B

INDEPENDENCE GROVE

LAKE FACTS

Municipality:

Libertyville

T44N, R11E, Section 4

Elevation:

640 feet

Major Watershed:

Des Plaines River

Sub-Watershed:

Upper Des Plaines

Surface Area:

129.3 acres

Shoreline Length:

4.8 miles

Maximum Depth:

52.0 feet

Average Depth:

15.0 feet

Lake Volume:

1944.66 acre-feet

Watershed Area:

335 acres

Lake Type:

Borrow Pit

Management Entities:

Lake County Forest Preserve
District

Current Uses:

Fishing, non-motorize boating,
swimming, and aesthetics.

mg/L, which was well below the county median of 7.8 mg/L. Low TSS values are typically correlated with good water clarity and can be beneficial to many aspects of the lake ecosystem, including the plant and fish communities. Water clarity was measured by Secchi disk depth with the lowest reading in September (7.6 ft) and the deepest in June (13.6 ft) corresponding to the highest and lowest TSS values. The average Secchi depth for the season was 10.31 feet, which is significantly greater than the county median (2.98 ft). The average conductivity reading for Independence Grove was 0.7526 mS/cm, which was approximately 4.6% lower than the county median (0.7889 mS/cm). The Secchi depth for 2016 (10.31 ft) has improved greatly since 2000 when the average was 5.82 feet, which was most likely due to it being newly developed. The increase in aquatic plant coverage along with Zebra Mussels in the lake may also contribute to low nutrient concentrations and greater clarity.

Phosphorus and nitrogen are two important nutrients for algal growth. The average total phosphorus for Independence Grove was <0.013 mg/L compared to the county median of 0.067 mg/L. Total Kjeldahl nitrogen (TKN) is a measure of organic nitrogen, and is typically bound up in algal and plant cells. The average TKN for Independence Grove was 0.57 mg/L and was lower than the county median of 2.43 mg/L.


The aquatic plant community was assessed in July when most of the plants are likely to be present. Seventy-two percent of the sites sampled had aquatic plants present with 20 different species and one macro-algae (*Chara* spp.) found in Independence Grove. Coontail was the most abundant species found at 44% of the sites followed by American Pondweed and Elodea occurring at 23% of the sites. Both, Eurasian Watermilfoil and Curlyleaf Pondweed (also present in the lake), are invasive, exotic species, which can crowd out native species and should be kept under control in order for native species to thrive.

An erosion assessment was conducted in 2016 to determine the condition of the shoreline at the water/land interface. Overall, 45% of the shoreline had no erosion, 43% had slight erosion, 10% had moderate, and 2% had severe erosion. The area of severe erosion should be addressed soon. Most of the erosion appeared to be where fishermen access the lake. Rock platforms for fishermen have been added around the lake for shore access. Some type of control should be employed in the other areas to allow access for fishing, but also protect the shoreline from further erosion. It is much easier and less costly to mitigate slightly eroding shorelines than those with more severe erosion.


INDEPENDENCE GROVE WATERSHED

The lake is located in the Upper Des Plaines sub basin, within the Des Plaines River watershed. A watershed is a drainage basin where water from rain or snow melt drains into a body of water, such as a river, lake, reservoir, wetland or storm drain. The source of a lakes water supply is very important in determining its water quality and choosing management practices to protect the lake. Independence Grove receives a majority of its water through inlets and runoff from the road and parking lots with in the surrounding area. The Village of Libertyville has banned the use of lawn fertilizer with phosphorus in March of 2015.

The Independence Grove watershed is approximately 1945 acres. The area around the lake is comprised mainly of water (37.9%), public and private open space (33.8%) and forest and grassland (16.1 %). Public and Private open space and contribute 46.4% and 36.3%, respectively of the estimated runoff in the watershed. Most of the watershed is with in the Lake County Forest Preserve Boundary. The retention time of the lake was calculated as 19.32 years.



INDEPENDENCE GROVE WATERSHED AND LAND USE



INDEPENDENCE GROVE LAND USE 2016

Land Use	Acreage	Runoff Coeff.	Estimated Runoff, acft.	% Total of Estimated Runoff
Agricultural	19.97	0.05	2.7	2.7
Forest and Grassland	53.96	0.05	7.4	7.4
Government and Institutional	5.06	0.50	7.0	6.9
Public and Private Open Space	113.22	0.15	46.7	46.4
Single Family	0.22	0.50	0.3	0.3
Transportation	15.63	0.85	36.5	36.3
Water	126.74	0.00	0.0	0.0
TOTAL	334.79		100.7	100.0

VLMP — WATER QUALITY

VOLUNTEER LAKE MONITOR PROGRAM

VOLUNTEERS MEASURE WATER CLARITY USING THE SECCHI DISK TWICE A MONTH MAY THROUGH OCTOBER. IN 2016 THERE WERE 52 LAKES PARTICIPATING IN LAKE COUNTY.

IF YOU WOULD LIKE MORE INFORMATION PLEASE CONTACT:

ALANA BARTOLAI

(847) 377-8009


ABARTOLAI2@LAKE COUNTYIL.GOV

WHAT YOU CAN DO TO IMPROVE WATER QUALITY ON INDEPENDENCE GROVE

- Do not throw leaves, grass clippings, pet waste, and other organic debris into the street or driveway. Runoff carries these through storm sewers, directly into the lake.
- Build a rain garden to filter run-off from roofs, parking lots, and streets. This allows the phosphorus to be bound to the soil so it does not reach surface waters.
- Plant a buffer around your lake shoreline to reduce runoff and filter nutrients from entering your lake.

The VLMP was established in 1981 by the Illinois Environmental Protection Agency (IEPA) to be able to collect information on Illinois inland lakes, and to provide an educational program for citizens. The volunteers are primarily lakeshore residents, lake owners/managers, members of environmental groups, and citizens with interest in a particular lake. The VLMP relies on volunteers to gather information on their chosen lake. The primary measurement by volunteers is Secchi depth (water clarity). Water clarity can provide an indication of the general water quality of the lake. Other observations such as water color, suspended algae and sediment, aquatic plants and odor are also recorded. The sampling season is May through October with measurements taken twice a month.

Participation provides annual data that helps document water quality impacts and support lake management decisions. The VLMP program has provides lakes with annual baseline data that can be used to determine long term water quality trends and support current lake management decision making. The volunteers will provide data that is vital for the management of this lake. If you would like to participate or need more information about becoming a VLMP please contact the LCHD-ES.



WATER CLARITY

Water clarity is an indicator of water quality related to chemical and physical properties. Measurements taken with a Secchi disk indicate the light penetration into a body of water. Algae, microscopic animals, water color, eroded soil, and resuspension of bottom sediment are factors that interfere with light penetration and reduce water transparency. If light penetration is reduced significantly, macrophyte growth may be decreased which would in turn impact the organisms dependent upon them for food and cover.

The 2016 average clarity for Independence Grove was 10.31 feet; this was a 10.7% decrease in the lakes transparency since 2007 of 11.54 feet and the water clarity was above the county median of 2.98 feet. The shallowest Secchi depth for Independence Grove was in September (7.6 ft) and the deepest was in June (13.6 ft). Overall, the good water clarity in Independence Grove can be attributed to a healthy plant population and a small watershed. Aquatic plants compete with algae for nutrients and stabilize bottom substrate, which in turn improves water clarity.


Secchi disk transparency is an indicator overall water quality. In general, the greater the Secchi disk depth, the clearer the water and better the water quality. High TSS values are typically correlated with poor water clarity. Over the past 16 years water clarity in Independence Grove has improved significantly.

TOTAL SUSPENDED SOLIDS

Another measure of water clarity is turbidity, which is caused by particles of matter rather than the dissolved organic compounds. Suspended particles dissipate light, which may limit the depth plants can grow. The total suspended solid (TSS) parameter (turbidity) is composed of nonvolatile suspended compounds (NVSS), non-organic clay or sediment materials, and volatile suspended solids (TVS) (algae and other organic matter).

Seasonal Secchi readings changes are affected by algal growth. The absence or low density of algae in early spring usually provides deeper clarity but as the water warms clarity decreases with more algae present in the water. High turbidity caused by sediment or algae can shade out native aquatic plants, resulting in their reduction or disappearance from the littoral zone. This eliminates the benefits provided by plants, such as habitat for many fish species and stabilization of the lake bottom. The 2016 TSS concentrations in Independence Grove averaged 1.6 mg/L which was below the county median of 7.8 mg/L and lower than the 2007 average concentration of 2.2 mg/L. Low TSS values are typically correlated with good water clarity (Secchi disk depth) and can be beneficial to many aspects of the lake ecosystem including the plant and fish communities. Lakes with NVSS values ≥ 12 mg/l could cause impairment for aquatic life in inland lakes.

There are internal and external sources of sediment affecting the turbidity in Independence Grove. Internal sources of sediment suspension include wind and wave action along with carp spawning and feeding activity especially along eroded shoreline. Aquatic plants act as a buffer by preventing the resuspension of sediments from affecting the rest of the lake. External sources include sediments that are transported into the lake from culverts, bank erosion and other sources in the watershed. The average calculated nonvolatile suspended solids (NVSS) was 1.01 mg/L. The low NVSS means that means that there is very little amount suspended solids found in the water column.



TSS Total Suspended Solids
TSS are particles of algae or sediment suspended in the water column.
TVS Total Volatile Solids
TVS represents the fraction of total solids that are organic in nature, such as algae cells
NVSS Non-Volatile Suspended Solids
NVSS represents the non-organic clay and sediments that are suspended in the water column.
TDS Total Dissolved Solids
TDS are the amount of dissolved substance such as salts or minerals in the water after evaporation.

**WHAT HAS BEEN DONE
TO REDUCE PHOSPHORUS
LEVELS IN INDEPENDENCE
GROVE**

March 2015— Libertyville becomes the 11th community in Lake County to ban the use of fertilizer containing phosphorus.

July 2010- The State of Illinois passed a law to reduce the amount of phosphorus content in dishwashing and laundry detergents.

July 2010- The State of Illinois passed another law restricting the use of lawn fertilizers containing phosphorus by commercial applicators.



SALTS DISSOLVE AND MOVE DOWNHILL OR INTO THE NEAREST STORM DRAIN WITH STORM-WATER AND SNOWMELT RUNOFF TO THE NEAREST LAKE, RIVER OR POND. THEY DO NOT SETTLE OUT; THEY REMAIN IN THE WATER CYCLE VIRTUALLY FOREVER.

NUTRIENTS

The nutrients organisms need to live or grow are typically taken in from the environment. In a lake the primary nutrients needed for aquatic plant and algal growth are phosphorus and nitrogen. In most lakes, including Independence Grove, phosphorus is the limiting nutrient, which means everything that plants and algae need to grow is available in excess: sunlight, warm temperature, and nitrogen.

Phosphorus has a direct effect on the amount of plant and algal growth in lakes. The 2016 average total phosphorus (TP) epilimnion (near surface sample) concentration in Independence Grove was 0.013 mg/L, this was slight decrease from the 2007 concentration (0.014 mg/L). Lakes with concentrations exceeding 0.050 mg/L can support high densities of algae and aquatic plants, which can reduce water clarity and dissolved oxygen levels and are considered impaired by the IEPA. Phosphorus originates from a variety of sources, many of which are related to human activities which include: human and animal waste, soil erosion, septic systems, common carp, and runoff from farmland and lawns. Independence Grove's healthy plant population, small water shed and lake origin as a borrow pit is responsible for keeping the TP level low which results in less algae blooms during the summer. Nitrogen is the other nutrient critical for algal growth. Total Kjeldahl nitrogen is a measure of organic nitrogen, and is typically bound up in algal and plant cells. The average 2016 TKN for Independence Grove was 0.47 mg/L. If inorganic nitrogen (NO₂, NO₃, NH₄) concentrations exceed 0.3 mg/L in spring, sufficient nitrogen is available to support summer algae blooms. However, low nitrogen levels do not guarantee less algae blooms. The TN:TP (total nitrogen to total phosphorus) ratio looks at which nutrient was limiting plant and algal growth in a lake. Ratios < 10:1 indicate nitrogen was limiting. Ratios >15:1 indicate phosphorus was limiting. Ratios >10:1, <15:1 indicate there was enough of both nutrients for excessive algal growth. The TN:TP ratio for Independence Grove was 42:1, which means that the limiting nutrient for aquatic plants was phosphorus.



CONDUCTIVITY AND CHLORIDE

Conductivity is a measure of a water's ability to conduct electricity, measured by the water's ionic activity and content. The higher the concentration of (dissolved) ions the higher the conductivity becomes. Conductivity readings, which are influenced by chloride concentrations, have been increasing throughout the past decade in Lake County. Lakes with residential and/or urban land uses in their watershed often have higher conductivity readings and higher Cl⁻ concentrations because of the use of road salts. Storm water run-off from impervious surfaces such as roads and parking lots can deliver high concentrations of Cl⁻ to nearby water bodies. Road salt used in the winter road maintenance consists of the following ions: sodium chloride, calcium chloride, potassium chloride, magnesium chloride, or ferrocyanides which are detected when chlorides are analyzed.

The 2016 average conductivity for Independence Grove was 0.7526 mS/cm. This parameter was lower the county median of 0.7889 mS/cm which is 15.2% lower than the 2007 value of 0.9308 mS/cm. These values are influenced by the winter road maintenance of Route 137, River Road and the surrounding roads and parking lots in the park. The United States Environmental Protection Agency has determined that chloride concentrations higher than 230 mg/L can disrupt aquatic systems and prolonged exposure can harm 10% of aquatic species. Independence Grove's Cl⁻ concentration was 83.9 mg/L which is 9% lower than the 2007 value of 92.3 mg/l. Chlorides tend to accumulate within a watershed as these ions do not break down and are not utilized by plants or animals. High chloride concentrations may make it difficult for many of our native species to survive. However, many of our invasive species, such as Eurasian Watermilfoil, Cattail and Common Reed, are tolerant to high chloride concentrations.

TROPHIC STATE INDEX

Another way to look at phosphorus levels and how they affect lake productivity is to use a Trophic State Index (TSI) based on phosphorus (TSIp). TSIp values are commonly used to classify and compare lake productivity levels (trophic state). A lake's response to additional phosphorus is an accelerated rate of eutrophication. Eutrophication is a natural process where lakes become increasingly enriched with nutrients. Lakes start out with clear water and few aquatic plants and over time become more enriched with nutrients and vegetation until the lake becomes a wetland. This process takes thousands of years to take place. Human activities on a lake or in the watershed accelerate this process by resulting in rapid soil erosion and heavy phosphorus inputs. This accelerated aging process on a lake is referred to as cultural eutrophication. The TSI index classifies the lake into one of four categories: oligotrophic (nutrient-poor, biologically unproductive), mesotrophic (intermediate nutrient availability and biological productivity), and eutrophic (nutrient rich, highly productive), or hypereutrophic (extremely nutrient-rich, productive). In 2016, Independence Grove was mesotrophic with a TSIp Value of 40.4, placing it 3rd out of 175 lakes in the county. Lake Carina and Sterling Lake was 1st with a 38.7 TSIp Value.



“WHEN HUMAN ACTIVITIES ACCELERATE LAKE EUTROPHICATION, IT IS REFERRED TO AS CULTURAL EUTROPHICATION. CULTURAL EUTROPHICATION MAY RESULT FROM SHORELINE EROSION, AGRICULTURAL AND URBAN RUNOFF, WASTEWATER DISCHARGES OR SEPTIC SEEPAGE, AND OTHER NON-POINT SOURCE POLLUTION SOURCES.”

LAKE LEVEL

Lakes with stable water levels potentially have less shoreline erosion problems. The lake level in Independence Grove was measured from a bridge at the entrance of North Bay. The lake level decreased from May to September by 15.6 inches. The highest water level recorded occurred in May (6.0 ft) and the lowest level in May (7.30 ft). The most significant water level fluctuation occurred from May to June with a decrease in the lake level of 10.8 inches. Independence Grove’s water level appear to be influenced by rain events. The drainage system at the park surrounding the lake contributes to the water level in the lake. In order to accurately monitor water levels it is recommended that a staff gauge be installed and levels measured and recorded frequently (daily or weekly). The data provides lake managers a much better idea of lake level fluctuations relative to rainfall events and can aid in future decisions regarding lake level. Staff gauge is a great tool for measuring water level in lakes, rivers, reservoirs. The data collected can be compiled to help understand the natural fluctuations of the lake. Lakes with fluctuating water levels potentially have poorer water quality and have more shoreline erosion problems.



EXAMPLE OF A PERMANENT STAFF GAUGE

2016	Level (ft)	Seasonal Change (ft)	Monthly Change (ft)
May	6.00		
June	6.90	0.90	0.90
July	7.06	1.06	0.16
August	7.30	1.30	0.24
September	6.83	0.83	-0.47

BATHYMETRIC MAPS

Bathymetric maps are also known as depth contour maps and display the shape and depth of a lake. They are valuable tools for lake managers because they provide information about the surface area and volume of the lake at certain depths. This information can then be used to determine the volume of lake that goes anoxic, how much of the lake bottom can be inhabited by plants, and is essential in the application of whole-lake herbicide treatments, harvesting activities and alum treatments of your lake. Other common uses for the map include sedimentation control, fish stocking, and habitat management.


STRATIFICATION

A lake's water quality and ability to support fish are affected by the extent to which the water mixes. The depth, size, and shape of a lake are the most important factors influencing mixing, but climate, lakeshore topography, inflow from streams and vegetation also play a role. Variations in density caused by different temperatures can prevent warm and cold water from mixing, called stratification.


For example: when lake ice melts in early spring, the temperature and density of lake water will be similar from top to bottom. Since it is uniform throughout the water column, the lake can mix completely recharging the bottom water with oxygen and bringing nutrients up to the surface. Some lakes in summer experience stratification where the lake is dividing into three zones: epilimnion (warm surface layer), thermocline (transition zone between warm and cold water) and hypolimnion (cold bottom water). Stratification traps nutrients released from bottom sediments in the hypolimnion and prevents mixing. The hypolimnion typically becomes anoxic (dissolved oxygen <1 mg/L).

Monthly depth profiles were measured on Independence Grove by measuring water temperature, dissolved oxygen, conductivity, and pH every foot from the lake surface to the lake bottom. The relative thermal resistance to mixing (RTRM) value can be calculated from this data which can tell us if the lake stratifies, how great the stratification is, and what depth it occurs.

Independence Grove began to stratify in mid-May and it was 14 feet in June and July, 16 feet in August and 20 feet in September as the water began to cool. The lake became anoxic at 42 feet in June and it lasted into September. This pattern of decreased oxygen content as the summer progresses is normal for the hypolimnion.



Independence Grove Temperature Profile 2016



BLUE-GREEN ALGAE

Algae are important to the freshwater ecosystems, and most species of algae are not harmful. Algae blooms are often caused by blue-green algae, or “cyanobacteria”, which are similar to bacteria in structure but utilize photosynthesis to grow. They have no nucleus and lack the photosynthetic pigments found in algae. They usually are too small to be seen individually, but can form visible colonies that can cover large areas of lakes. Certain species of blue-green algae can produce toxins that could pose a health risk to people and animals when they are exposed to them in large enough quantities.

Independence Grove did not have a seasonal algae bloom due to a healthy aquatic plant population that competed with algae for nutrients and stabilize bottom substrate, which in turn improves water clarity. This allowed the total phosphorus levels to remain low at an average of 0.016 mg/l during the season.

When a bloom occurs, the water can appear blue-green, bright green, brown, or red and may look like paint floating on the water. Not all blue-green algae produce harmful toxins. The three types of cyanobacteria that are often associated with Harmful Algal Bloom (HAB) are the Anabaena, Aphanizomenon, and Microcystis. The presence of these cyanobacteria does not generally mean that the toxins are present in the water. The presence of toxins can only be verified through a sample analyzed in the lab. Poisoning has caused the death of cows, dogs, and other animals. Most human cases occurred when people swim or ski in affected recreational water bodies during a bloom.

If you suspect that you are experiencing symptoms related to exposure to blue-green algae such as stomach cramps, diarrhea, vomiting, headache, fever, muscle weakness, or difficulty breathing contact your doctor or the poison control center. For more information or to report a blue-green algae bloom, contact the Lake County Health Department Environmental Services (847) 377-8030.

FOR MORE INFORMATION
ON BLUE-GREEN ALGAE:
www.epa.state.il.us/

[water/surface-water/blue-green-algae.html](http://www.epa.state.il.us/water/surface-water/blue-green-algae.html)


TO REPORT BLUE-GREEN
ALGAE BLOOM:
Lake County Health
Department
847-377-8030



Anabaena Sp.



Microcystis Sp.



Aphanizomenon Sp.



FLORISTIC QUALITY INDEX

LAKE COUNTY
AVERAGE
FQI = 13.4

INDEPENDENCE GROVE
FQI = 24.4

RANK = 12/171

AQUATIC PLANTS
SPECIES
OBSERVED = 20

Floristic quality index (FQI) is an assessment tool designed to evaluate the closeness that the flora of an area is to that of undisturbed conditions. It can be used to: 1) identify natural areas, 2) compare the quality of different sites or different locations within a single site, 3) monitor long-term floristic trends, and 4) monitor habitat restoration efforts. Each aquatic plant in a lake is assigned a number between 1 and 10 (10 indicating the plant species most sensitive to disturbance). This is done for every floating and submersed plant species found in the lake. These numbers are averaged and multiplied by the square root of the number of species present to calculate an FQI. A high FQI number indicates that there are a large number of sensitive, high quality plant species present in the lake. Non-native species were counted in the FQI calculations for Lake County lakes. In 2016, Independence Grove had an FQI of 24.4 ranking 12 out of 171 in Lake County. The median FQI of lakes that we have studied from 2000-2016 is 13.4. Cedar Lake is 1st with an FQI of 37.4.

In many lakes macrophytes contribute to the aesthetically pleasing appearance of the setting and are enjoyable in their own right. but even more important, they are an essential element in the life systems of most lakes.

**AQUATIC PLANTS:
WHERE DO THEY GROW?**



Littoral Zone– the area that aquatic plants grow in a lake.

Algae– have no true roots, stems, or leaves and range in size from tiny, one-celled organisms to large, multicelled plant-like organisms.

Submerged Plants– have stems and leaves that grow entirely underwater, although some may also have floating leaves.

Floating-leaf Plants– are often rooted in the lake bottom, but their leaves and flowers float on the water surface.

Emergent Plants– are rooted in the lake bottom, but their leaves and stems extend out of the water.



← **Littoral Zone** →

Source: Minnesota Department of Natural Resources


AQUATIC PLANTS

Aquatic plant mapping survey provides information based on the species, density and distribution of plant communities in a particular lake. An aquatic plant sampling was conducted on Independence Grove on July 2016 when most plants are present. There were 147 points generated based on a computer grid system with points 60 meters apart. Aquatic plants occurred at 106 of the sites (72% total lake coverage) that included 20 aquatic plant species, including 2 exotic invasive species: Curlyleaf Pondweed and Eurasian Watermilfoil. The macro algae Chara, was also found during the plant survey. The most commonly occurring species was Coontail (44.2%), American Pondweed (23.1%), Elodea (23.1%), Water Stargrass (20.4%), Eurasian Watermilfoil (19.7%), and White Water Crowfoot (17%) were the next abundant species. Northern and Brittle Naiad are 2 new aquatic plant observed in 2016 while Spiny Naiad and Leafy pondweed were not observed during the survey. The extent of plant populations can be influenced by a variety of factors. Water clarity and depth are the major limiting factors in determining the maximum depth at which aquatic plants will grow. When light level in the water column falls below 1% of the surface light level, plants can no longer grow. The extent of the 1% light can be obtained by doubling the Secchi disk reading. The average Secchi disk reading for 2016 was 10.31 feet. The deepest aquatic plant, Coontail, was found in 15 feet of water. Aquatic plants play an important role in the lakes ecosystem by providing habitat for fish and shelter for aquatic organism. Aquatic plants provide oxygen, reduce nutrients such as phosphorus to prevent algae bloom, and help stabilize sediment.



LCHD Staff identifying plants during sampling.

AQUATIC PLANT MAP FOR JULY 2016



Rake Density (coverage)	# of Sites	% of Sites
No Plants	26	31
>0-10%	19	23
10-40%	18	21
40-60%	10	12
60-90%	2	2
>90%	9	11
Total Sites with Plants	58	69
Total # of Sites	84	100

AQUATIC PLANTS



Chara is an advance form of algae which resemble higher plants. Its easily identified by its musky odor and gritty surface due to mineral deposits on its surface. It filters nutrients out of the water and stabilizes the lake bottom.

© Paul Skawinski, 2009



Water Stargrass



Sago Pondweed



Paul Skawinski, 2009

Illinois Pondweed

Aquatic plants provide many water quality benefits and play an important role in the lakes ecosystem by providing habitat for fish and shelter for aquatic organism. Plants provide oxygen, reduce nutrients such as phosphorus to prevent algae bloom, and help stabilize sediment. A native plant community tends to be diverse and usually does not impede lake activities such as boating, swimming and fishing.


COMMON AQUATIC PLANTS FOUND IN INDEPENDENCE GROVE 2016

COONTAIL (*Ceratophyllum demersum*)



This perennial plant is a submerged aquatic about 1-3' long. There is more branching of the stems above than below, creating fan-like aggregations of leaves. The stems are up to 1.0 mm. across, light green to nearly white, terete to slightly compressed (flattened), and hairless; they are slender and flexible. The leaves are highly flexible and readily bend. The preference is full sun, shallow water up to 4' deep, and a mucky bottom.

JULY 2016 COONTAIL DENSITY AT INDEPENDENCE GROVE



AQUATIC PLANTS

AMERICAN PONDWEED (*Potamogeton nodosus*)



American pondweed is a perennial plant that has both floating and a few submerged leaves in an alternate pattern. The floating leaves are elliptical to oval 4 to 7 inches long and to 2 1 inches wide on long petioles. Submerged leaves are not abundant and are blade-like, somewhat transparent and smaller than floating leaves. Rhizomes are white. Fruits are on spikes that often stand above the water's surface and are brownish to reddish, 1 inches long and 1/8 to 1/4 inches wide.



Southern Naiad

ELODEA (*Elodea canadensis*)



Elodea canadensis, commonly called Canadian pondweed, is a submerged aquatic perennial. It produces fragile, branching, tangled stems to 3.5' long (10-12' in the wild) densely clad with short, drooping, lance-shaped to ovate dark green leaves (to 5/8" long) in whorls of three. Plants may grow completely submersed and at some considerable depth, or in shallower water the growing tips of stems may float on the surface. It produces tiny greenish-white flowers appear from July to September.



Slender Naiad


WHITE WATER CROWFOOT (*Ranunculus aquatilis*)



A single flower on a long naked stalk arising from a leaf axil. Flowers are 1/3 to 2/3 inch across, have 5 oval white petals with a spot of yellow at the base, and many yellow stamens surrounding a yellow center that matures to green. 5 rounded green sepals behind the flower are much shorter than the petals.

Leaves are finely divided into many thread-like segments, round to fan-shaped in outline, to 3/4 inch long and about 1 inch wide, alternately attached. Stems and leaves are all submersed and mostly hairless.

The flower center expands to a round to oval seed head, the seeds smooth to roughly hairy and with a beak almost as long as the seed body.




White Water Lily


EURASIAN WATERMILFOIL

ILLUSTRATION OF EURASIAN WATERMILFOIL

Illustration provided by:
IFAS, Center for Aquatic Plants
University of Florida, Gainesville, 1990



ERASIAN WATERMILFOIL DENSITY AT 29 SITES ON INDEPENDENCE GROVE IN JULY, 2016



Eurasian Watermilfoil (EWM) is a feathery submerged aquatic plant that can quickly form thick mats in shallow areas of lakes and rivers in North America. These mats can interfere with swimming and entangle propellers, which hinders boating fishing, and waterfowl hunting. Matted milfoil can displace native aquatic plants, impacting fish and wildlife. Since it was discovered in North America in the 1940's, EWM has invaded nearly every US state and at least three Canadian Provinces. Milfoil spreads when plant pieces break off and float on water currents. It can cross land to new waters by clinging to sailboats, personal watercraft, powerboats, motors, trailers, and fishing gear.

The occurrence of EWM in Independence Grove has decreased from 2014 (35.6%) to 2016



(19.7%) of the sample sites. An aquatic plant management plan is critical to maintaining the health of the lake and a balanced aquatic plant community. The plan should be based on the management goals of the lake and involve usage issues, habitat maintenance/restoration, and limitations of the lake. The primary focus of the plan must include the control of exotic aquatic species including EWM and Curlyleaf Pondweed. Follow up is critical to achieve long-term success. A good aquatic plant management plan considers both the short and long-term needs of the lake. At this time there is a healthy population of native aquatic plants to keep the Eurasian water milfoil from becoming the dominant plant in the lake. Hand raking areas may help reduce the EWM population.



MYRIOPHYLLUM SPICATUM **EXOTIC***

COMMON NAMES:
EURASIAN WATERMILFOIL

ORIGIN: EXOTIC
EUROPE AND ASIA. FOUND THROUGHOUT LAKE COUNTY AND ILLINOIS

IMPORTANCE:
THIS INVASIVE PLANT SPREADS RAPIDLY, CROWDING OUT NATIVE SPECIES, CLOGGING WATERWAYS, AND BLOCKING SUNLIGHT AND OXYGEN FROM UNDERLYING WATERS.

LOOK ALIKES:
NORTHERN WATERMILFOIL WHICH HAS FEWER THAN 12 LEAFLET PAIRS PER LEAF, AND GENERALLY HAS STOUTER STEMS.



KEY FEATURES:

STEM: LONG, OFTEN ABUNDANTLY BRANCHED STEMS FORM A REDDISH OR OLIVE-GREEN SURFACE MAT IN SUMMER.

LEAF: LEAVES ARE RECTANGULAR WITH ≥12 PAIRS OF LEAFLETS PER LEAF AND ARE DISSECTED GIVING A FEATHERY APPEARANCE, ARRANGED IN A WHORL, WHORLS ARE 1 INCH APART.


FLOWER: SMALL PINKISH MALE FLOWERS THAT OCCUR ON REDDISH SPIKES, FEMALE FLOWERS LACK PETALS AND SEPALS AND 4 LOBED PISTIL.

CURLYLEAF PONDWEED


Curlyleaf Pondweed (CLP) is an invasive aquatic perennial that is native to Eurasia, Africa, and Australia. It was accidentally introduced to United States waters in the mid-1880s by hobbyists who used it as an aquarium plant. The leaves are reddish-green, oblong, and about 3 inches long, with distinct wavy edges that are finely toothed. The stem of the plant is flat, reddish-brown and grows from 1 to 3 feet long. This aquatic plant has an unusual life history. Unlike our native pondweeds it begins growing in the early spring. CLP has even been documented growing under the ice in lakes! The plant then reaches maturity in mid summer typically June in Lake County when our natives are starting to emerge. CLP becomes invasive in some areas because of its adaptations for low light tolerance and low water temperatures which allow the plant to get a head start and outcompete native plants in the spring. In mid-summer, when most aquatic plants are growing, CLP plants are dying off. Plant die-offs may result in a critical loss of dissolved oxygen. Large populations of CLP also can cause changes in nutrient availability. In midsummer, CLP plants usually die back which is typically followed by an increase in phosphorus availability that may fuel nuisance algal blooms. CLP can form dense mats that may interfere with boating and other recreational uses. In July 2016, Independence Grove CLP were present, plants being found at 4.1% of the sampled sites. There were more CLP observed in May and June especially in North Bay. At this time the density of CLP is not causing fluctuations in nutrient availability. The LCFP should monitor the CLP and manually remove the plants using a hand rake to keep the population from expanding.



ILLUSTRATION OF CURLYLEAF PONDWEED




CURLYLEAF PONDWEED DENSITY AT 6 SITES ON INDEPENDENCE GROVE IN JULY 2016



POTAMOGETON CRISPUS **EXOTIC***



Turion



KEY FEATURES:

STEM: ARE FLATTENED, BRANCHED, CAN FORM DENSE STANDS IN WATER UP TO 15 FEET DEEP.

LEAF: ALTERNATE ALL SUBMERSED, OBLONG, STIFF, TRANSLUCENT LEAVES HAVE DISTINCTLY WAVY EDGES WITH FINE TEETH AND 3 MAIN VEINS.

FLOWER: TINY, WITH 4 PETAL-LIKE LOBES. IN SPIKES 1-3CM LONG ON STALKS UP TO 7CM LONG. (MAY SEE TURIONS WHICH OVER WINTERS AS A HARD, BROWN, BUR-LIKE BUD WITH CROWDED, SMALL HOLLY-LIKE LEAVES).

COMMON NAMES:

CURLY LEAF PONDWEED

ORIGIN: EXOTIC*

ASIA, AFRICA, AND EUROPE FOUND THROUGHOUT LAKE COUNTY AND ILLINOIS

IMPORTANCE:

INVASIVE: HAS A TOLERANCE FOR LOW LIGHT AND WATER TEMPERATURES THAT ALLOW THE PLANT TO GET A HEAD START ON NATIVE PLANTS. BY MID SUMMER WHEN MOST AQUATIC PLANTS ARE GROWING, CURLYLEAF PLANTS ARE DYING OFF. WHICH MAY RESULT IN A CRITICAL LOSS OF DISSOLVED OXYGEN AND AN INCREASE IN NUTRIENTS.

LOOK ALIKES:

NONE

AQUATIC HERBICIDE 2-4-D



Copper Sulfate Application

Chemical applications for algae is a temporary solution that often requires multiple applications. As the treated algae sink to the bottom to decompose (use oxygen) they release nutrients that the surviving algae uses to rebound.

FOR FULL DETAILS OF THE RULE SEE:

[HTTP://
WWW.EPA.STATE.IL.
US/WATER/PERMITS/
PESTICIDE/
INDEX.HTML](http://www.epa.state.il.us/water/permits/pesticide/index.html)



Aquatic herbicide 2,4-D can be applied to the water in either a liquid or pellet form. 2,4-D is a relatively fast-acting, systemic, selective herbicide used for the control of Eurasian watermilfoil and other broad-leaved species. Both the granular and liquid formulations can be effective for spot treatment of Eurasian watermilfoil. 2,4-D has been shown to be selective to Eurasian watermilfoil when used at the labeled rate, leaving many native aquatic species relatively unaffected. 2,4-D herbicides will affect only dicots, broad leafed plants like Eurasian watermilfoil. Most native aquatic plants in our lakes are monocots. Proper planning should be taken when applying herbicides. If large quantities of aquatic plants are killed, their decomposition can reduce the amount of dissolved oxygen that can lead to fish kill. Rain, wind direction, boating activity, and water temperature may affect the effectiveness of the applied chemical.

COARSE WOODY HABITAT

The vegetation around Independence Grove is made up of native plants and shrubs. This results in the lack of coarse woody habitat (CWH) that is important to a lake's ecosystems. CWH often happens naturally as trees or large branches fall into the lake become material critical habitat for tiny aquatic organisms that feed fish, turtles, birds, and other wildlife. Sunken logs or fish habitat can be found in the lake along with CWH at the outlet bay. The rest of the lake does not have CWH along the shoreline.

Bluegills and young of the year game fish often seek shelter in and around CWH. Larger logs with branching limbs provide shade and protection for the smaller fish. Black Bass species (smallmouth and largemouth) often build spawning nests in proximity to CWH, particularly large logs (Hunt and Annett 2002; Lawson et al. 2011; Weis and Sass 2011).

Adding CWH in the littoral or near-shore zone serves many functions within a lake ecosystem including erosion control, as a carbon source, and as a surface for algal growth which is an important food base for aquatic macroinvertebrates (Engel and Pederson 1998; Sass 2009). Macroinvertebrates provide food for panfish and young of the year gamefish. Logs and branches that are out of the water are used by turtles, birds and other wildlife.



MANUAL REMOVAL OF AQUATIC PLANTS

Controlling exotic aquatic plants by hand removal is effective on small areas and if done prior to heavy infestation. Eurasian Watermilfoil can be controlled to some degree by hand pulling or raking of entire plants including the roots. Just before the peak growth is the best time for removal to prevent re-growth and plant seed dispersal. Working in windblown areas will help contain fragments near shore which makes cleanup easier. All fragments of EWM plants must be removed to achieve adequate control. Most regeneration are from fragmented stems that drift into different areas of the lake and form new colonies. Small populations of Curlyleaf can be manually removed with a rake or by hand pulling. All plant materials should be carefully removed as floating pieces of Curlyleaf that may contain turions can float away and colonize a different part of the lake so removal of this plant should be done no later that the end of May.

Removal by hand is labor intensive but it can be effective in targeting small patches of invasive plants. This method also eliminates or reduces the need for chemicals treatments that can impact native vegetation and fish. There are different types of rakes. First is a bladed rake that can be used to cut the stems of plants. Secondly, a throw able double sided rake that can be used to pull plants from deeper water or further distances and lastly a long handled rake for working the shoreline and the boat dock area. Its important to remove the entire plant including the roots to prevent regeneration.



Aquatic Rake with Blade

PROTECT YOUR WATERS

Helping to prevent the introduction and spread of invasive aquatic plants and animals are the most effective way of protecting healthy, non-infested ecosystems. Listed below are some of the simple steps you can take to prevent invasion.

- Remove all plants, mud, fish, or animals before transporting equipment.
- Eliminate all water from equipment before transporting equipment.
- Dry anything that comes in contact with water (boat, trailers, equipment, clothing, etc.).
- Remove all mud and dirt since it might contain aquatic hitchhikers.
- Never release plants, fish or animals into a body of water unless they came out of that body of water.
- Do not release bait into the waters you are fishing.
- Do not release aquarium fish or aquatic pets in to the lake.

STOP AQUATIC HITCHHIKERS!™
 Prevent the transport of nuisance species.
 Clean all recreational equipment.
www.ProtectYourWaters.net



Double-sided Throwable Rake





Aquatic Plant Rake

DISSOLVED OXYGEN

Dissolved oxygen (DO) is a major indicator of water quality and is important for aquatic organisms, algae, macrophytes, and for many chemical reactions to occur that are crucial for lake functions. Dissolved oxygen concentrations can have large variations occurring and are affected by diffusion, aeration, photosynthesis, respiration, and decomposition. Temperature, salinity and pressure changes will also cause DO to fluctuate. Dissolved oxygen will vary both seasonally and by depth throughout the water column in lakes. If dissolved oxygen concentrations drops below levels necessary for sustaining aquatic life (below 5.0 mg/L at 1 foot depth below the lake surface) it becomes a water quality impairment. Low dissolved oxygen primarily is a result of excessive nutrients that stimulate growth of organic matter, such as algae, or the increase of pollutants such as sewage, lawn clippings, and soils that are considered to be “oxygen-demanding”. Low dissolved oxygen levels is also often a factor for fish kills. When many of the plants or algae die at the end of the growing season, their decomposition can significantly reduce DO concentrations. In deeper, thermally stratified lakes, oxygen production is greatest in the upper water layer (epilimnion) where sunlight drives photosynthesis and oxygen consumption is greatest near the bottom of the lake (hypolimnion) where organic matter accumulates and decomposes.

Independence Grove’s dissolved oxygen concentrations remained good all year and did not drop below 5.0 mg/L at 1 foot below the lakes surface, which would qualify it as a DO impairment. The lake averaged 9.0 mg/L at 3 feet below the surface from May to September.

RANGE OF TOLERANCE FOR DISSOLVED OXYGEN IN FISH PARTS PER MILLION (PPM) DISSOLVED OXYGEN	
1	
2	<3.0 PPM too low for fish populations
3	
4	3.0-5.0 PPM 12-24 Hour range of tolerance /stressful conditions
5	
6	>6.0 PPM Supports Spawning
7	
8	>7.0 PPM Supports Growth and Activity
9	
10	>9.0 PPM Supports abundant fish populations



AQUATIC PLANTS AND FISH

Fish depend on aquatic plants to provide habitat and forage for food. Most freshwater fish rely on aquatic plants at some point during their life stage. The plant composition and density can play an important role in the nesting, growth, and foraging success of these fish. While many fish require some aquatic vegetation for growth, excessive amounts of aquatic vegetation can negatively impact growth by reducing foraging success. The parameters of an ideal fish habitat can change based on the size and species of fish, the type of lake, structures present in the lake and many other factors. A fish survey can reveal the size distribution and population of the fish in your lake.

How do plants impact fish?

- ◆ *Plants provide critical structure to aquatic habitats.*
- ◆ *Plants influence growth of fish by enhancing fish diversity, feeding, growth, and reproduction.*
- ◆ *Plants influence spawning. The structure provided by plant beds are important to fish reproduction.*
- ◆ *Plants influence the physical environment. Aquatic plants can change water temperatures and available oxygen in habitats.*




Image <http://agriculture.vic.gov.au/>

Table 1. Common fish and their plant affinity during various life stages and their relationship with plants

Fish	Plant Affinity	Life Stage				Relationship	
		Larvae	Juvenile	Adult	Spawn	Forage	Predator avoidance
Bluegill sunfish	High	X	X	X	X	X	X
Common carp	High	X	X	X	X	X	X
Largemouth bass	High	X	X	X	X	X	X
Musky	High	X	X	X	X	X	X
Northern Pike	High	X	X	X	X	X	X
Black crappie	Moderate		X	X	X	X	X
Smallmouth bass	Moderate		X	X		X	X
Yellow perch	Moderate	X	X			X	X
White crappie	Low		X			X	
Salmon, trout	Low		X				X
Shad	Low	X					
Walleye	Low			X		X	

Table adapted from Gettys, Lynn, William T. Haller and Marc Bellaud. "Biology and Control of Aquatic Plants: A Best Management Practices Handbook". 2009

SHORELINE EROSION

“VEGETATIVE BUFFER ZONES CAN PLAY A KEY ROLE IN LIMITING NEGATIVE WATER QUALITY IMPACTS FROM DEVELOPED SHORELAND PROPERTY”

Erosion is a natural process primarily caused by water which results in the loss of material from the shoreline. Disturbed shorelines caused by human activity such as clearing of vegetation and beach rocks, and increasing runoff will accelerate erosion. Rain and melting snow and wave action are the main causes of erosion. Rain can loosen soil and wash it down gradient towards the lake. Creating a native plant buffer helps prevent soil erosion as well as filter out pollutants and unwanted nutrients from entering the lake. Native plants can be planted along the shoreline since plant roots hold the soil particles in place so they are not easily washed away during a rain event, melting snow or wave action. Loose rocks and gravel placed on top of a filter fabric prevents soil from washing away before newly planted seed and vegetation has a chance to grow. Eroded materials cause turbidity, sedimentation, nutrients, and pollutants to enter a lake. Shore line buffer zone planted with native vegetation not only reduces runoff by increasing water infiltration into the ground, it also offers food and habitat for wildlife. Less runoff means less nutrients, sediments and other pollutants entering the lakes and streams. Excess nutrients are the primary cause of algal blooms and increased aquatic plant growth. Once in the lake, sediments, nutrients and pollutants are harder and more expensive to remove.

A shoreline erosion study was assessed for Independence Grove in 2016. The lake was divided into reaches, and the shoreline evaluated for none, slight, moderate and severe erosion based on exposed soil and tree/plant roots, failing infrastructure, undercut banks, and other signs of erosion. Based on the 2016 data, 55% of Independence Grove’s shoreline has some erosion; 43% slight erosion, 10% moderate erosion, and 2% severe erosion.



SHORELINE BUFFER

INDEPENDENCE GROVE’S SHORELINE EROSION

INFORMATION ON SHORELINE REGULATION AND PERMITS CAN BE FOUND ON THE ILLINOIS DEPARTMENT OF NATURAL RESOURCES’ WEBSITE.

HTTP://WWW.DNR.ILLINOIS.GOV/WATERRESOURCES/DOCUMENTS/3704.PDF



ZEBRA MUSSELS *DREISSENA POLYMORPHA*

In the late 1990's, the presence of zebra mussels (*Dreissena polymorpha*) was confirmed in the Fox Chain O Lakes. These mussels are believed to have been spread to this country in the mid 1980's by cargo ships from Europe that discharged their ballast water into the Great Lakes. The mussels spread throughout the Great Lakes and by 1991 had made their way into the Illinois and Mississippi Rivers. The first sighting of the mussel in Lake County (besides Lake Michigan and the Chain of Lakes) occurred in 1999. Currently, 33 inland lakes in the county are known to be infested with the zebra mussel, but this number could be much higher, since the zebra mussel has probably gone unnoticed in many lakes. Zebra mussels were first discovered in Independence Grove in 1999 and are now spread through out the entire lake.

The zebra mussel's reproductive cycle allows for rapid expansion of the population. A mature female can produce up to 40,000 eggs in a cycle and up to one million in a season. Zebra mussels can live as long as five years and have an average life span of about 3.5 years. The adults are typically about the size of a thumbnail but can grow as large as 2 inches in diameter. Colonies can reach densities of 30,000 - 70,000 mussels per square meter. Due to their quick life cycle and explosive growth rate, zebra mussels can quickly edge out native mussel species. Negative impacts on native bivalve populations include interference with feeding, habitat, growth, movement and reproduction. The impact that mussels have on fish populations is not fully understood. However, zebra mussels feed on phytoplankton (algae), which is also a major food source for planktivorous fish, such as minnows and shad and young of the year bluegill. These fish, in turn, are a food source for piscivorous fish (fish eating fish), such as largemouth bass and northern pike.

Zebra mussels have also caused economic problems for large power plants, public water supplies, and industrial facilities, where they clog water intake pipes. Boats stored on the water offer suitable areas for zebra mussels to start a colony. Researchers found that many of the mussel larvae were being transported via aquatic plants that were taken from one lake to another on boats and trailers. It is important that all boats and trailers entering or leaving Independence Grove are inspected for aquatic plants, zebra mussels and all water from the bilge and motors are drained.

Below are some tips from the Great Lakes Sea Grant Network that can help prevent the spread of zebra mussels:

- Always inspect your boat and boat trailer carefully before transporting. Studies have shown that transport via aquatic plant fragments is one of the major contributors to the spread of zebra mussels.
- Drain all bilge waters, live wells, bait buckets and engine compartments before entering another lake. Make sure water is not trapped in your trailer. Never transport water from one lake to another.
- Flush clean water (tap) through the cooling system of your motor to rinse out any larvae. Full grown zebra mussels can be easily seen but cling stubbornly to surfaces. Boats that have been in the water for long periods of time should be carefully inspected. Carefully scrape the hull (or trailer), or use a high pressure spray (250 psi) to dislodge them. Or leave your boat out of the water for at least 5 days, preferably up to two weeks. The mussels will die and drop off.
- In their earlier stages, attached zebra mussels may not be easily seen. Pass your hand across the boat's bottom - if it feels grainy, it's probably covered with mussels. Don't take a chance; clean them off by scraping or blasting.
- Dispose of the mussels in a trash barrel or other garbage container. Don't leave them on the shore where they could be swept back into the lake or foul the area.




ZEBRA MUSSEL

For more information:

[http://
www.seagrant.wisc.edu/
zebramussels/faqs.html](http://www.seagrant.wisc.edu/zebramussels/faqs.html)



ZEBRA MUSSEL
FORM A COLONY
CALLED "DRUSES"



ZEBRA ON A
ROCKY SHORELINE
DURING LOW
WATER

SWIMMING BEACH MONITORING

All licensed inland beaches are tested bi-weekly from May to September by the Lake County Health Department's Ecological Services Department. The water samples are tested for E. coli bacteria, which are found in the intestines of almost all warm-blooded animals. E. coli is used as an indicator organism, meaning that high concentrations of E. coli might suggest the presence of harmful pathogens such as, Salmonella, Giardia, etc. While not all strains of E. coli are the same, certain strains can make humans sick if ingested in high enough concentrations. If water samples come back high for E. coli (>235 E. coli/100 ml), LCHD informs the management body for the bathing beach that the beach is closed and a sign is posted indicating the beach closure. Licensed beaches will also be monitored for blue-green algae and samples may be taken for testing when blooms occur. During the summer of 2016, E. coli counts at Independence Grove's beach had one beach closure on August 23 due to high E. coli (410.6 E. coli/100ml). There were 45 water fowl observed at the beach when the sample was taken which could be the cause of the high E. coli count.

If you have any questions, feel free to contact the Lake County Health Department at 847-377-8030.

How to prevent illness and beach closure:

- If you are sick, do NOT swim.
- Don't swim when you have diarrhea. You can spread germs in the water.
- Take a shower prior to entering the beach area.
- Children who are not toilet trained should wear tight fitting rubber or plastic pants.
- Pick up garbage around the beach area.
- Avoid swimming during algae blooms.
- Do not ingest the water while swimming.

SWIMMING BEACH




LAKE MANAGEMENT PLANS

It is recommended that a long term Lake Management Plans be developed to effectively manage lake issues. All stakeholders should participate in the development of the plan and include homeowners, recreational users, lake management associations, park districts, townships or any other entity involved in managing Independence Grove. Lake Management plans should educate the public about specific lake issues, provide a concise assessment of the problem, outline methods and techniques that will be employed to control the problems and clearly define the goals of the program. Mechanisms for monitoring and evaluation should be developed as well and information gathered during these efforts should be used to implement management efforts (Biology and Control of Aquatic Plants, Gettys et al., 2009)

What are the steps in creating a Lake Management Plan?

1. **Getting Started:** Identify lake stakeholders and communication pathways
2. **Setting Goals:** Getting the effort organized, identifying problems to be addressed, and agreeing on the goals
3. **Problem Assessment & Analysis:** collecting baseline information to define the past and existing conditions. Synthesize the information, quantifying and comparing the current conditions to desired conditions, researching opportunities and constraints and setting direction to achieve goals.
4. **Alternatives:** List all possible management alternatives and evaluate their strengths, weakness, and general feasibility.
5. **Recommendations:** Prioritize management options, setting objectives and drafting the plan
6. **Project Management:** Management of assets, detailed records of expenses and time
7. **Implementation:** adopting the plan, lining up funding, and scheduling activities for taking action to achieve goals.
8. **Monitor & Modify:** Develop a mechanism for tracking activities and adjusting the plan as it evolves.

Follow these steps when getting started with writing Lake Management Plans. While each step is necessary, the level of effort and detail for each step will vary depending on the project's goals, size of the lake, and number of stakeholders.





ECOLOGICAL SERVICES

Senior Biologist: Mike Adam

madam@lakecountyiil.gov

Population Health Services
500 W. Winchester Road

Phone: 847-377-8030
Fax: 847-984-5622

For more information visit us at:

**[http://www.lakecountyiil.gov/
Health/want/
BeachLakeInfo.htm](http://www.lakecountyiil.gov/Health/want/BeachLakeInfo.htm)**

Protecting the quality of our lakes is an increasing concern of Lake County residents. Each lake is a valuable resource that must be properly managed if it is to be enjoyed by future generations. To assist with this endeavor, Population Health Environmental Services provides technical expertise essential to the management and protection of Lake County surface waters.

Ecological Service's goal is to monitor the quality of the county's surface water in order to:

- Maintain or improve water quality and alleviate nuisance conditions
- Promote healthy and safe lake conditions
- Protect and improve ecological diversity

Services provided are either of a technical or educational nature and are provided by a professional staff of scientists to government agencies (county, township and municipal), lake property owners' associations and private individuals on all bodies of water within Lake County.

LAKE RECOMMENDATIONS


The water quality and aquatic plant coverage has improved in Independence Grove since 2007. With a reduction in total suspended solids (27%), total phosphorus (7%) and chloride (9%). Since 2014 aquatic plant diversity increased from 14 aquatic plants to 20 aquatic plants observed in 2016. Independence Grove's management is administered by the Lake County Forest Preserve (LCFP).

To improve the overall quality of Independence Grove, ES (Ecological Services) has the following recommendations:


- **Participation in Volunteer Lake Monitoring Program**
- **Install a sign to educate on ways to reduce the spread of Aquatic Invasive Species**
- **Help reduce Cl⁻ by supporting wise use of road salt in the watershed**
- **Install a permanent staff gage to monitor lake level fluctuations**
- **Become familiar with the appearance of harmful algal blooms and report any blooms to the LCHD-ES by calling 847-837-8030**
- **Develop an Aquatic Plant Management Plan (APMP) that targets the reduction of invasive species and promotes native plant diversity. Aquatic plant management plans should consider type, timing of pesticide applications and quantity of pesticide used**
- **Monitor Eurasian Watermilfoil and Curlyleaf; population. Hand rake or manually remove Eurasian Watermilfoil and Curlyleaf to keep population from expanding**
- **Incorporate native plants in their landscaping through rain gardens shoreline buffer**
- **Mitigate shoreline exhibiting erosion**
- **Asses current fish population and add course woody habitat (CWD) to improve the lake's ecosystem**

Appendix A:
Figures



Independence Grove 2016 Sampling Point





Independence Grove Plant Grid 2016





Independence Grove Aquatic Plant Biovolume July 2016



Independence Grove Coontail Rake Density July 2016




Independence Grove EWM Rake Density July 2016




Independence Grove Watershed



 Independence Grove Watershed




Independence Grove Land Use 2016




- Agricultural
- Forest and Grassland
- Government and Institutional
- Public and Private Open Space
- Single Family
- Transportation
- Water
- IndependenceGroveShed07


Indipendence Grove Shoreline Buffer 2016



— Good
— Fair
— Poor





Indipendence Grove Shoreline Buffer 2016





	Good
	Fair
	Poor

Independence Grove Shoreline Erosion 2016



Independence Grove Shoreline Reaches, 2016



Appendix B:
Tables

Lake County average Floristic Quality Index (FQI) ranking 2000-2016.

RANK	LAKE NAME	FQI (w/A)	FQI (native)
1	Cedar Lake	37.4	38.9
2	East Loon Lake	34.7	36.1
3	Cranberry Lake	29.7	29.7
4	Deep Lake	29.7	31.2
5	Round Lake Marsh North	29.1	29.9
6	West Loon Lake	27.1	29.5
7	Sullivan Lake	26.9	28.5
8	Bangs Lake	26.2	27.8
9	Little Silver Lake	25.2	26.7
10	Third Lake	25.1	22.5
11	Fourth Lake	24.7	27.1
12	Indpendence Grove	24.4	26.8
13	Sun Lake	24.3	26.1
14	Redwing Slough	24.0	25.8
15	Schreiber Lake	23.9	24.8
16	Lakewood Marsh	23.8	24.7
17	Sterling Lake	23.6	25.4
18	Deer Lake	23.5	24.4
19	Round Lake	23.5	25.9
20	Pistakee Lake	23.5	25.2
21	Lake Marie	23.5	25.2
22	Lake of the Hollow	23.0	24.8
23	Nippersink Lake (Fox Chain)	22.4	23.2
24	Countryside Glen Lake	21.9	22.8
25	Grass Lake	21.5	22.2
26	Davis Lake	21.4	21.4
27	Timber Lake (North)	20.9	23.4
28	Lake Catherine	20.8	21.8
29	Cross Lake	20.7	18.7
30	ADID 203	20.5	20.5
31	Broberg Marsh	20.5	21.4
32	McGreal Lake	20.2	22.1
33	Fox Lake	20.2	21.2
34	Honey Lake	20.0	20.0
35	Lake Barrington	19.9	21.8
36	Lake Kathryn	19.6	20.7
37	Druce Lake	19.1	21.8
38	Turner Lake	18.6	21.2
39	Salem Lake	18.5	20.2
40	Duck Lake	18.3	19.2
41	Wooster Lake	18.0	20.1
42	Lake Helen	18.0	18.0
43	Old Oak Lake	18.0	19.1
44	Lake Minear	18.0	20.1
45	Potomac Lake	17.8	17.8
46	Lake Zurich	17.7	18.9
47	Redhead Lake	17.7	18.7
48	Long Lake	17.7	15.8
49	Hendrick Lake	17.7	17.7
50	Rollins Savannah 2	17.7	17.7
51	Grandwood Park Lake	17.2	19.0
52	Seven Acre Lake	17.0	15.5
53	Lake Miltmore	16.8	18.7
54	Petite Lake	16.8	18.7
55	Channel Lake	16.8	18.7
56	McDonald Lake 1	16.7	17.7
57	Highland Lake	16.7	18.9
58	Almond Marsh	16.3	17.3

Lake County average Floristic Quality Index (FQI) ranking 2000-2016.

RANK	LAKE NAME	FQI (w/A)	FQI (native)
59	Owens Lake	16.3	17.3
60	Windward Lake	16.3	17.6
61	Butler Lake	16.1	18.1
62	Grays Lake	16.1	16.1
63	White Lake	16.0	17.0
64	Dunns Lake	15.9	17.0
65	Dog Bone Lake	15.7	15.7
66	Osprey Lake	15.5	17.3
67	Heron Pond	15.1	15.1
68	Ames Pit	15.1	17.6
69	North Churchill Lake	15.0	15.0
70	Hastings Lake	15.0	17.0
71	Forest Lake	14.8	15.9
72	Dog Training Pond	14.7	15.9
73	Summerhill Estates Lake	14.5	15.5
74	Grand Ave Marsh	14.3	16.3
75	Nippersink Lake	14.3	16.3
76	Taylor Lake	14.3	16.3
77	Manning's Slough	14.1	16.3
78	Tower Lake	14.0	14.0
79	Dugdale Lake	14.0	15.1
80	Eagle Lake (S1)	14.0	15.1
81	Crooked Lake	14.0	16.0
82	Spring Lake	14.0	15.2
83	Lake Matthews	13.9	15.5
84	Longview Meadow Lake	13.9	13.9
85	Fischer Lake	13.6	14.7
86	Bishop Lake	13.4	15.0
87	Mary Lee Lake	13.1	15.1
88	Old School Lake	13.1	15.1
89	Lake Tranquility (S1)	12.6	12.6
90	Buffalo Creek Reservoir 1	12.5	11.4
91	Buffalo Creek Reservoir 2	12.5	11.4
92	McDonald Lake 2	12.5	12.5
93	Rollins Savannah 1	12.5	12.5
94	Stone Quarry Lake	12.5	12.5
95	Kemper Lake 1	12.2	13.4
96	Pond-A-Rudy	12.1	12.1
97	Stockholm Lake	12.1	13.5
98	Lake Leo	12.1	14.3
99	Lambs Farm Lake	12.1	14.3
100	Bresen Lake	12.0	13.9
101	Grassy Lake	12.0	12.0
102	Flint Lake Outlet	11.8	13.0
103	Albert Lake	11.5	10.3
104	Rivershire Pond 2	11.5	13.3
105	Hook Lake	11.3	13.4
106	Briarcrest Pond	11.2	12.5
107	Lake Naomi	11.2	12.5
108	Pulaski Pond	11.2	12.5
109	Lake Napa Suwe	11.0	11.0
110	Redwing Marsh	11.0	11.0
111	West Meadow Lake	11.0	11.0
112	Nielsen Pond	10.7	12.0
113	Lake Holloway	10.6	10.6
114	Sylvan Lake	10.6	10.6
115	Echo Lake	10.4	10.4
116	Gages Lake	10.2	12.5

Lake County average Floristic Quality Index (FQI) ranking 2000-2016.

RANK	LAKE NAME	FQI (w/A)	FQI (native)
117	College Trail Lake	10.0	10.0
118	Valley Lake	9.9	9.9
119	Werhane Lake	9.8	12.0
120	Fish Lake	9.6	10.6
121	Lake Carina	9.5	12.5
122	Columbus Park Lake	9.2	9.2
123	Lake Lakeland Estates	9.2	9.2
124	Waterford Lake	9.2	9.2
125	Bluff Lake	9.1	11.0
126	Lake Fairfield	9.0	10.4
127	Des Plaines Lake	8.6	9.9
128	Antioch Lake	8.5	8.5
129	Loch Lomond	8.5	8.5
130	Lake Fairview	8.5	6.9
131	Timber Lake (South)	8.5	6.9
132	East Meadow Lake	8.5	8.5
133	South Churchill Lake	8.5	8.5
134	Kemper Lake 2	8.5	9.8
135	Lake Christa	8.5	9.8
136	Lake Farmington	8.5	9.8
137	Lucy Lake	8.5	9.8
138	Lake Louise	8.4	8.4
139	Bittersweet Golf Course #13	8.1	8.1
140	Lake Linden	8.0	8.0
141	Sand Lake	8.0	10.4
142	Countryside Lake	7.7	11.5
143	Fairfield Marsh	7.5	8.7
144	Lake Eleanor	7.5	8.7
145	Banana Pond	7.5	9.2
146	Slocum Lake	7.1	5.8
147	Lucky Lake	7.0	7.0
148	North Tower Lake	7.0	7.0
149	Lake Forest Pond	6.9	8.5
150	Ozaukee Lake	6.7	8.7
151	Leisure Lake	6.4	9.0
152	Peterson Pond	6.0	8.5
153	Little Bear Lake	5.8	7.5
154	Deer Lake Meadow Lake	5.2	6.4
155	ADID 127	5.0	5.0
156	Island Lake	5.0	5.0
157	Liberty Lake	5.0	5.0
158	Oak Hills Lake	5.0	5.0
159	Slough Lake	5.0	5.0
160	International Mining and Chemical Lake	5.0	7.1
161	Lochanora Lake	5.0	5.0
162	Diamond Lake	3.7	5.5
163	Lake Charles	3.7	5.5
164	Big Bear Lake	3.5	5.0
165	Sand Pond (IDNR)	3.5	5.0
166	Harvey Lake	3.3	5.0
167	Half Day Pit	2.9	5.0
168	Hidden Lake	0.0	0.0
169	St. Mary's Lake	0.0	0.0
170	Willow Lake	0.0	0.0
171	Woodland Lake	0.0	0.0
	<i>Mean</i>	13.9	14.9
	<i>Median</i>	13.4	14.7

Lake County average TSI phosphorus (TSIp) ranking 2000-2016.

RANK	LAKE NAME	TP AVE	TSIp
1	Sterling Lake	0.0110	38.73
2	Lake Carina	0.0110	38.73
3	Independence Grove	0.0130	40.38
4	Cedar Lake	0.0130	41.14
5	Druce Lake	0.0140	42.21
6	Windward Lake	0.0160	44.13
7	Lake Minear	0.0164	44.49
8	Sand Pond (IDNR)	0.0165	44.57
9	West Loon	0.0170	45.00
10	Pulaski Pond	0.0180	45.83
11	Ames Pit	0.0190	46.61
12	Banana Pond	0.0200	47.35
13	Gages Lake	0.0200	47.35
14	Lake Kathryn	0.0200	47.35
15	Highland Lake	0.0202	47.49
16	Lake Miltmore	0.0210	48.05
17	Timber Lake (North)	0.0210	48.05
18	Lake Zurich	0.0210	48.05
19	Cross Lake	0.0216	48.46
20	Dog Training Pond	0.0220	48.72
21	Sun Lake	0.0220	48.72
22	Deep Lake	0.0230	49.36
23	Lake of the Hollow	0.0230	49.36
24	Round Lake	0.0230	49.36
25	Stone Quarry Lake	0.0230	49.36
26	Bangs Lake	0.0260	51.13
27	Lake Leo	0.0260	51.13
28	Lake Barrington	0.0270	51.68
29	Cranberry Lake	0.0270	51.68
30	Dugdale Lake	0.0270	51.68
31	Peterson Pond	0.0270	51.68
32	Little Silver Lake	0.0280	52.20
33	Wooster Lake	0.0290	52.71
34	Lambs Farm Lake	0.0310	53.67
35	Old School Lake	0.0310	53.67
36	Grays Lake	0.0310	53.67
37	Harvey Lake	0.0320	54.13
38	Butler Lake	0.0324	54.31
39	Hendrick Lake	0.0340	55.00
40	Fourth Lake	0.0360	55.82
41	Sullivan Lake	0.0370	56.22
42	Sand Lake	0.0380	56.60
43	Third Lake	0.0384	56.77
44	Diamond Lake	0.0390	56.98
45	East Loon	0.0400	57.34
46	Schreiber Lake	0.0400	57.34
47	Waterford Lake	0.0400	57.34
48	Hook Lake	0.0410	57.70
49	Lake Tranquility (S1)	0.0412	57.77
50	Nielsen Pond	0.0450	59.04
51	Seven Acre Lake	0.0460	59.36

Lake County average TSI phosphorus (TSIp) ranking 2000-2016.

RANK	LAKE NAME	TP AVE	TSIp
52	Turner Lake	0.0460	59.36
53	Willow Lake	0.0460	59.36
54	East Meadow Lake	0.0480	59.97
55	Lucky Lake	0.0480	59.97
56	Old Oak Lake	0.0490	60.27
57	College Trail Lake	0.0500	60.56
58	Summerhill Estates Lake	0.0514	60.96
59	Hastings Lake	0.0520	61.13
60	West Meadow Lake	0.0530	61.40
61	Lucy Lake	0.0550	61.94
62	Lake Linden	0.0570	62.45
63	Lake Christa	0.0580	62.70
64	Owens Lake	0.0580	62.70
65	Briarcrest Pond	0.0580	62.70
66	Honey Lake	0.0586	62.85
67	Redhead Lake	0.0608	63.38
68	St. Mary's Lake	0.0608	63.38
69	Duck Lake	0.0610	63.43
70	Lake Lakeland Estates	0.0620	63.66
71	Lake Naomi	0.0620	63.66
72	Lake Catherine	0.0620	63.66
73	Liberty Lake	0.0630	63.89
74	North Tower Lake	0.0630	63.89
75	Werhane Lake	0.0630	63.89
76	Countryside Glen Lake	0.0640	64.12
77	Davis Lake	0.0650	64.34
78	Leisure Lake	0.0650	64.34
79	Channel Lake	0.0680	65.00
80	Buffalo Creek Reservoir 1	0.0680	65.00
81	Mary Lee Lake	0.0680	65.00
82	Little Bear Lake	0.0680	65.00
83	Crooked Lake	0.0710	65.62
84	Timber Lake (South)	0.0720	65.82
85	Lake Helen	0.0720	65.82
86	Grandwood Park Lake	0.0720	65.82
87	ADID 203	0.0730	66.02
88	Fish Lake	0.0730	66.02
89	Broberg Marsh	0.0780	66.97
90	Echo Lake	0.0790	67.16
91	Countryside Lake	0.0800	67.34
92	Lake Nippersink	0.0800	67.34
93	Woodland Lake	0.0800	67.34
94	Redwing Slough	0.0822	67.73
95	Tower Lake	0.0830	67.87
96	Antioch Lake	0.0850	68.21
97	Potomac Lake	0.0850	68.21
98	White Lake	0.0862	68.42
99	Grand Ave Marsh	0.0870	68.55
100	North Churchill Lake	0.0870	68.55
101	McDonald Lake 1	0.0880	68.71
102	Pistakee Lake	0.0880	68.71

Lake County average TSI phosphorus (TSIp) ranking 2000-2016.

RANK	LAKE NAME	TP AVE	TSIp
103	Lake Fairview	0.0890	68.88
104	Rivershire Pond 2	0.0900	69.04
105	South Churchill Lake	0.0900	69.04
106	McGreal Lake	0.0910	69.20
107	Lake Charles	0.0930	69.51
108	Deer Lake	0.0940	69.66
109	Eagle Lake (S1)	0.0950	69.82
110	International Mine and Chemical Lake	0.0950	69.82
111	Valley Lake	0.0950	69.82
112	Buffalo Creek Reservoir 2	0.0960	69.97
113	Big Bear Lake	0.0960	69.97
114	Fox Lake	0.1000	70.56
115	Nippersink Lake - LCFP	0.1000	70.56
116	Sylvan Lake	0.1000	70.56
117	Petite Lake	0.1020	70.84
118	Longview Meadow Lake	0.1020	70.84
119	Lake Marie	0.1030	70.98
120	Dunn's Lake	0.1070	71.53
121	Lake Forest Pond	0.1070	71.53
122	Long Lake	0.1070	71.53
123	Grass Lake	0.1090	71.80
124	Des Plaines Lake	0.1090	71.80
125	Spring Lake	0.1100	71.93
126	Kemper 2	0.1100	71.93
127	Bittersweet Golf Course #13	0.1100	71.93
128	Osprey Lake	0.1110	72.06
129	Bluff Lake	0.1120	72.19
130	Middlefork Savannah Outlet 1	0.1120	72.19
131	Lochanora Lake	0.1120	72.19
132	Round Lake Marsh North	0.1130	72.32
133	Deer Lake Meadow Lake	0.1160	72.70
134	Lake Matthews	0.1180	72.94
135	Taylor Lake	0.1180	72.94
136	Island Lake	0.1210	73.31
137	Columbus Park Lake	0.1230	73.54
138	Lake Holloway	0.1320	74.56
139	Fischer Lake	0.1380	75.20
140	Slocum Lake	0.1500	76.40
141	Lakewood Marsh	0.1510	76.50
142	Pond-A-Rudy	0.1510	76.50
143	Forest Lake	0.1540	76.78
144	Bresen Lake	0.1580	77.15
145	Middlefork Savannah Outlet 2	0.1590	77.24
146	Grassy Lake	0.1610	77.42
147	Salem Lake	0.1650	77.78
148	Half Day Pit	0.1690	78.12
149	Lake Louise	0.1810	79.11
150	Lake Eleanor	0.1810	79.11
151	Lake Farmington	0.1850	79.43
152	ADID 127	0.1890	79.74

Lake County average TSI phosphorus (TSIp) ranking 2000-2016.

RANK	LAKE NAME	TP AVE	TSIp
153	Lake Napa Suwe	0.1940	80.11
154	Loch Lomond	0.1960	80.26
155	Patski Pond	0.1970	80.33
156	Dog Bone Lake	0.1990	80.48
157	Redwing Marsh	0.2070	81.05
158	Stockholm Lake	0.2082	81.13
159	Bishop Lake	0.2160	81.66
160	Ozaukee Lake	0.2200	81.93
161	Kemper 1	0.2220	82.06
162	Hidden Lake	0.2240	82.19
163	McDonald Lake 2	0.2250	82.25
164	Oak Hills Lake	0.2790	85.35
165	Heron Pond	0.2990	86.35
166	Rollins Savannah 1	0.3070	86.73
167	Fairfield Marsh	0.3260	87.60
168	ADID 182	0.3280	87.69
169	Manning's Slough	0.3820	89.88
170	Slough Lake	0.3860	90.03
171	Rasmussen Lake	0.4860	93.36
172	Albert Lake, Site II, outflow	0.4950	93.62
173	Flint Lake Outlet	0.5000	93.76
174	Rollins Savannah 2	0.5870	96.08
175	Almond Marsh	1.9510	113.40
	<i>Average</i>	<i>0.1113</i>	<i>65.8</i>

Lake County Secchi Disk Clarity Ranking, 2000-2016.

RANK	LAKE NAME	SECCHI AVE	TSI_{sd}
1	Lake Carina	16.96	36.31
2	Windward Lake	14.28	38.79
3	Sterling Lake	13.84	39.24
4	Cedar Lake	12.55	40.66
5	Druce Lake	12.25	41.00
6	Pulaski Pond	11.69	41.68
7	West Loon Lake	11.55	41.85
8	Lake Zurich	10.40	43.37
9	Independence Grove	10.31	43.49
10	Ames Pit	9.97	43.97
11	Third Lake	9.76	44.28
12	Davis Lake	9.65	44.44
13	Harvey Lake	9.47	44.72
14	Little Silver Lake	9.42	44.79
15	Old School Lake	9.40	44.82
16	Lake Kathryn	9.39	44.84
17	Dugdale Lake	9.22	45.10
18	Dog Training Pond	9.04	45.39
19	Banana Pond	8.85	45.69
20	Deep Lake	8.83	45.72
21	Stone Quarry Lake	8.81	45.76
22	Wooster Lake	8.74	45.87
23	Lake of the Hollow	8.74	45.87
24	Cross Lake	8.18	46.83
25	Bangs Lake	8.02	47.11
26	Briarcrest Pond	8.00	47.15
27	Sand Lake	7.48	48.12
28	Sand Pond (IDNR)	7.42	48.23
29	Timber Lake (North)	7.37	48.33
30	Lake Miltmore	7.35	48.37
31	Lake Leo	7.31	48.45
32	Schreiber Lake	7.25	48.57
33	Nielsen Pond	7.23	48.61
34	Honey Lake	7.17	48.73
35	Lake Minear	7.13	48.81
36	Round Lake	7.01	49.05
37	Highland Lake	6.97	49.14
38	Lake Helen	6.43	50.30
39	Sun Lake	6.33	50.52
40	Lake Barrington	6.12	51.01
41	Cranberry Lake	5.94	51.44
42	Lake Fairfield	5.89	51.56
43	Gages Lake	5.45	52.68
44	Owens Lake	5.30	53.08
45	Valley Lake	5.05	53.78
46	McGreal Lake	5.04	53.81
47	Old Oak Lake	4.85	54.36
48	Waterford Lake	4.70	54.82
49	Lake Linden	4.60	55.13
50	Timber Lake (South)	4.46	55.57
51	Peterson Pond	4.51	55.41

Lake County Secchi Disk Clarity Ranking, 2000-2016.

RANK	LAKE NAME	SECCHI AVE	TSIsd
52	Crooked Lake	4.39	55.79
53	Mary Lee Lake	4.35	55.93
54	Butler Lake	4.35	55.93
55	Crooked Lake	4.28	56.17
56	Deer Lake	4.20	56.45
57	Seven Acre Lake	4.18	56.51
58	Lambs Farm Lake	4.17	56.54
59	Grays Lake	4.08	56.86
60	Lake Naomi	4.05	56.96
61	White Lake	3.96	57.29
62	Hook Lake	3.95	57.32
63	Turner Lake	3.92	57.43
64	Leisure Lake	3.85	57.69
65	Summerhill Estates Lake	3.84	57.73
66	North Tower Lake	3.89	57.74
67	Salem Lake	3.77	58.00
68	Lake Fariview	3.75	58.07
69	Duck Lake	3.71	58.23
70	Countryside Glen Lake	3.64	58.50
71	Fish Lake	3.57	58.78
72	Lochanora	3.52	58.99
73	Taylor Lake	3.52	58.99
74	Hastings Lake	3.52	58.99
75	Bishop Lake	3.47	59.19
76	Lake Lakeland Estates	3.41	59.44
77	Lake Holloway	3.40	59.49
78	Stockholm Lake	3.38	59.57
79	East Loon Lake	3.30	59.92
80	Lucky Lake	3.22	60.27
81	Diamond Lake	3.17	60.50
82	Liberty Lake	3.16	60.54
83	International Mining and Chemical Lake	3.08	60.91
84	Long Lake	3.05	61.05
85	Lake Christa	3.01	61.24
86	Lucy Lake	2.99	61.34
87	Lake Catherine	2.9	61.78
88	St. Mary's Lake	2.79	62.34
89	Channel Lake	2.77	62.44
90	Werhane Lake	2.71	62.76
91	Bresen Lake	2.69	62.86
92	East Meadow Lake	2.61	63.30
93	Buffalo Creek Reservoir 1	2.60	63.35
94	Countryside Lake	2.58	63.46
95	Kemper Lake 1	2.56	63.58
96	Bluff Lake	2.51	63.86
97	Broberg Marsh	2.50	63.92
98	Antioch Lake	2.48	64.03
99	Little Bear Lake	2.38	64.63
100	Island Lake	2.32	65.00
101	Tower Lake	2.31	65.06
102	Buffalo Creek Reservoir 2	2.30	65.12

Lake County Secchi Disk Clarity Ranking, 2000-2016.

RANK	LAKE NAME	SECCHI AVE	TSIsd
103	Woodland Lake	2.28	65.25
104	Rivershire Pond 2	2.23	65.57
105	Lake Charles	2.20	65.76
106	Fischer Lake	2.70	62.81
107	College Trail Lake	2.18	65.89
108	Loch Lomond	2.17	65.96
109	Redhead Lake	2.16	66.03
110	Pistakee Lake	2.15	66.09
111	Echo Lake	2.11	66.36
112	Eagle Lake (S1)	2.10	66.43
113	West Meadow Lake	2.07	66.64
114	Forest Lake	2.04	66.85
115	Grand Ave Marsh	2.03	66.92
116	Columbus Park Lake	2.03	66.92
117	Grassy Lake	2.00	67.14
118	Petite Lake	2	67.14
119	Des Plaines Lake	2.14	66.16
120	Sylvan Lake	1.98	67.28
121	Bittersweet Golf Course #13	1.98	67.28
122	Spring Lake	1.78	68.82
123	Kemper Lake 2	1.77	68.90
124	Fourth Lake	1.77	68.90
125	Nippersink Lake	1.73	69.23
126	Deer Lake Meadow Lake	1.73	69.23
127	Lake Louise	1.68	69.65
128	Willow Lake	1.63	70.09
129	Slough Lake	1.63	70.09
130	Rasmussen Lake	1.62	70.17
131	Lake Farmington	1.62	70.17
132	Half Day Pit	1.60	70.35
133	Lake Marie	1.56	70.72
134	Longview Meadow Lake	1.51	71.19
135	Lake Matthews	1.48	71.48
136	Big Bear Lake	1.32	73.13
137	Fox Lake	1.28	73.57
138	Dunn's Lake	1.22	74.26
139	Lake Eleanor	1.16	74.99
140	McDonald Lake 1	1.13	75.37
141	Lake Napa Suwe	1.06	76.29
142	Rollins Savannah 1	1.05	76.43
143	Osprey Lake	1.03	76.70
144	Manning's Slough	1.00	77.13
145	Rollins Savannah 2	0.95	77.87
146	Dog Bone Lake	0.94	78.02
147	Redwing Marsh	0.88	78.97
148	Flint Lake Outlet	0.83	79.82
149	Fairfield Marsh	0.81	80.17
150	Slocum Lake	0.81	80.17
151	Oak Hills Lake	0.79	80.53
152	Grass Lake	0.78	80.71
153	Lake Nippersink	0.77	80.90

Lake County Secchi Disk Clarity Ranking, 2000-2016.

RANK	LAKE NAME	SECCHI AVE	TSIsd
154	South Churchill Lake	0.73	81.67
155	Lake Forest Pond	0.71	82.07
156	ADID 127	0.66	83.12
157	North Churchill Lake	0.61	84.26
158	Hidden Lake	0.56	85.54
159	Ozaukee Lake	0.51	86.84
160	McDonald Lake 2	0.50	87.12

Independence Grove Water Quality Summary 2016

2016		Eplimnion														
DATE	DEPTH	ALK	TKN	NH ₃ -N	NO ₂ +NO ₃ -N	TP	SRP	TDS	Cl	TSS	TS	TVS	SECCHI	COND	pH	DO
17-May	3	165	0.45	<0.1	<0.05	<0.01	<0.005	437	81	1.7	527	128	8.75	0.7750	8.28	10.8
14-Jun	3	147	0.48	<0.1	<0.05	0.01	<0.005	425	84	0	545	165	13.6	0.7500	8.17	9.4
12-Jul	3	135	0.45	<0.1	<0.05	<0.01	<0.005	428	86.5	2.1	545	181	10.00	0.7570	8.28	8.9
16-Aug	3	129	0.45	<0.1	<0.05	0.02	<0.005	424	84.0	2.2	495	134	11.60	0.7480	8.2	8.0
13-Sep	3	127	0.52	<0.1	<0.05	0.01	<0.005	416	84	2.2	505	145	7.6	0.7330	8.33	8.2

Average 141 0.57 <0.1^k <0.05 0.013^k <0.005 426 83.9 1.6 523 151 10.31 0.7526 8.3 9.0

2007		Eplimnion														
DATE	DEPTH	ALK	TKN	NH ₃ -N	NO ₂ +NO ₃ -N	TP	SRP	TDS	Cl	TSS	TS	TVS	SECCHI	COND	pH	DO
15-May	3	139	0.55	<0.1	<0.05	0.016	<0.005	NA	92.2	2.7	618	140	9.19	0.9610	8.34	9.6
19-Jun	3	124	0.56	<0.1	<0.05	<0.01	<0.005	NA	92.0	1.8	648	198	11.80	0.9180	8.43	8.6
17-Jul	3	126	0.60	<0.1	<0.05	<0.01	<0.005	NA	94.5	2.1	606	128	14.10	0.9480	8.55	8.0
14-Aug	3	120	0.57	<0.1	<0.05	<0.01	<0.005	NA	92.5	1.6	599	150	13.12	0.9280	8.17	7.0
18-Sep	3	125	<0.5	<0.1	<0.05	0.011	<0.005	NA	90.5	2.6	622	177	9.51	0.8990	8.56	11.1

Average 127 0.57^k <0.1 <0.05 0.014^k <0.005 NA 92.3 2.2 619 159 11.54 0.9308 8.4 8.9

2000		Eplimnion														
DATE	DEPTH	ALK	TKN	NH ₃ -N	NO ₂ -N [*]	TP	SRP	TDS	Cl	TSS	TS	TVS	SECCHI	COND	pH	DO
10-May	3	210	0.67	<0.1	0.065	0.012	<0.005	712	NA	2.5	724	170	6.79	1.0870	8.62	8.3
14-Jun	3	197	1.34	<0.1	0.053	0.020	0.007	658	NA	3.8	817	291	5.80	1.0550	8.15	7.2
11-Jul	3	191	0.69	<0.1	<0.05	0.024	<0.005	696	NA	4.2	715	197	5.40	1.0020	8.11	6.9
16-Aug	3	197	0.53	<0.1	<0.05	0.021	<0.005	650	NA	3.6	751	229	5.41	1.0330	8.15	6.5
13-Sep	3	196	0.75	<0.1	<0.05	0.020	<0.005	626	NA	3.6	672	162	5.71	1.0420	8.03	7.3

Average 198 0.80 <0.1 0.059^k 0.019 0.007^k 668 NA 3.5 736 210 5.82 1.0440 8.21 7.2

Glossary
ALK = Alkalinity, mg/L CaCO ₃
TKN = Total Kjeldahl nitrogen, mg/L
NH ₃ -N = Ammonia nitrogen, mg/L
NO ₂ +NO ₃ -N = Nitrate + Nitrite nitrogen, mg/L
NO ₃ -N = Nitrate nitrogen, mg/L
TP = Total phosphorus, mg/L
SRP = Soluble reactive phosphorus, mg/L
Cl = Chloride, mg/L
TDS = Total dissolved solids, mg/L
TSS = Total suspended solids, mg/L
TS = Total solids, mg/L
TVS = Total volatile solids, mg/L
SECCHI = Secchi disk depth, ft.
COND = Conductivity, milliSiemens/cm
DO = Dissolved oxygen, mg/L

k = Denotes that the actual value is known to be less than the value presented.

NA= Not applicable

* = Prior to 2006 only Nitrate - nitrogen was analyzed

2016		Hypolimnion															
DATE	DEPTH	ALK	TKN	NH ₃ -N	NO ₂ +NO ₃ -N	TP	SRP	TDS	Cl	TSS	TS	TVS	SECCHI	COND	pH	DO	
17-May	42	247	1.12	0.607	<0.05	0.014	<0.005	557	85.7	3.1	760	193	NA	1.0130	7.43	2.06	
14-Jun	40	273	1.06	0.654	<0.05	0.021	0.006	628	89.0	1.7	862	239	NA	1.154	7.36	1.26	
12-Jul	36	265	0.92	0.49	<0.05	0.027	<0.005	626	87.4	6.3	867	278	NA	1.1510	7.32	0.67	
16-Aug	41	314	1.37	0.920	<0.05	0.029	<0.005	733	94.0	1.9	987	291	NA	1.364	7.28	0.33	
13-Sep	34	269	0.94	<0.01	0.48	0.0	<0.005	642	89.0	4.1	858	260	NA	1.1820	7.31	0.23	

Average 274 1.08 0.668^k 0.48^k 0.022 0.006^k 637 89.0 3.4 867 252 NA 1.1728 7.34 0.9

2007		Hypolimnion															
DATE	DEPTH	ALK	TKN	NH ₃ -N	NO ₂ +NO ₃ -N	TP	SRP	TDS	Cl	TSS	TS	TVS	SECCHI	COND	pH	DO	
15-May	42	194	0.64	0.245	0.089	<0.01	<0.005	NA	84.0	2.8	943	187	NA	1.3570	7.20	4.53	
19-Jun	38	243	0.88	0.491	<0.05	0.012	<0.005	NA	78.0	4.1	1330	301	NA	1.5530	7.10	0.80	
17-Jul	43	270	1.31	0.892	<0.05	0.035	<0.005	NA	75.6	13.0	1360	229	NA	1.7535	7.70	0.33	
14-Aug	45	294	1.72	1.330	<0.05	0.114	0.021	NA	76.0	20.6	1450	280	NA	1.7825	6.83	0.09	
18-Sep	42	299	1.70	1.360	<0.05	0.129	0.058	NA	75.4	19.0	1460	233	NA	1.8280	7.90	0.07	

Average 260 1.25 0.864 0.089^k 0.073 0.040^k NA 77.8 11.9 1309 246 NA 1.6548 7.35 1.2

2000		Hypolimnion															
DATE	DEPTH	ALK	TKN	NH ₃ -N	NO ₂ -N*	TP	SRP	TDS	Cl	TSS	TS	TVS	SECCHI	COND	pH	DO	
10-May	44	237	1.30	0.672	0.061	0.026	0.01	716	NA	3	758	174	NA	1.121	8.04	1.7	
14-Jun	40	238	0.91	0.776	0.071	0.030	0.013	749	NA	6	863	289	NA	1.131	7.5	0	
11-Jul	43	250	NA	1.100	<0.05	0.117	0.098	808	NA	3.7	863	272	NA	1.125	7.29	0	
16-Aug	44	273	2.00	1.780	<0.05	0.277	0.234	712	NA	2.5	773	179	NA	1.123	7.45	0	
13-Sep	43	277	2.60	1.970	<0.05	0.306	0.272	702	NA	3.6	748	196	NA	1.137	7.43	0.5	

Average 255 1.36 1.26 0.066^k 0.151 0.1254 737 NA 3.8 801 222 NA 1.127 7.54 0.4

Glossary
ALK = Alkalinity, mg/L CaCO ₃
TKN = Total Kjeldahl nitrogen, mg/L
NH ₃ -N = Ammonia nitrogen, mg/L
NO ₂ +NO ₃ -N = Nitrate + Nitrite nitrogen, mg/L
NO ₃ -N = Nitrate nitrogen, mg/L
TP = Total phosphorus, mg/L
SRP = Soluble reactive phosphorus, mg/L
Cl = Chloride, mg/L
TDS = Total dissolved solids, mg/L
TSS = Total suspended solids, mg/L
TS = Total solids, mg/L
TVS = Total volatile solids, mg/L
SECCHI = Secchi disk depth, ft.
COND = Conductivity, milliSiemens/cm
DO = Dissolved oxygen, mg/L

k = Denotes that the actual value is known to be less than the value presented.

NA= Not applicable

* = Prior to 2006 only Nitrate - nitrogen was analyzed

Independence Grove Multi-Parameter 2016

Date	Depth	Dep25	Temp	DO	DO%	SpCond	pH	BG
MMDDYY	feet	feet	°C	mg/l	Sat	mS/cm	Units	
5/17/2016	0	0.500	12.39	10.75	100.8	0.774	8.56	285
5/17/2016	1	1.076	12.39	10.76	100.9	0.774	8.46	221
5/17/2016	2	2.142	12.40	10.75	100.9	0.774	8.39	972
5/17/2016	3	3.028	12.40	10.75	100.9	0.775	8.28	558
5/17/2016	4	4.073	12.40	10.74	100.8	0.775	8.22	289
5/17/2016	6	6.048	12.39	10.73	100.7	0.774	8.20	684
5/17/2016	8	8.019	12.39	10.72	100.6	0.774	8.17	460
5/17/2016	10	10.090	12.34	10.70	100.3	0.774	8.16	711
5/17/2016	12	12.022	12.65	10.62	99.8	0.774	8.15	196
5/17/2016	14	14.027	12.29	10.52	98.5	0.774	8.11	708
5/17/2016	16	16.179	12.21	10.50	98.1	0.775	8.10	484
5/17/2016	18	18.096	11.86	10.44	96.6	0.777	8.09	474
5/17/2016	20	20.040	11.18	10.05	91.6	0.778	8.07	89
5/17/2016	22	22.040	7.42	9.90	82.5	0.796	8.03	379
5/17/2016	24	24.124	6.66	8.75	71.5	0.805	7.96	361
5/17/2016	26	26.025	6.44	8.23	67.1	0.811	7.88	899
5/17/2016	28	28.125	6.24	7.81	62.8	0.817	7.82	1393
5/17/2016	30	30.120	6.19	7.35	59.2	0.830	7.78	450
5/17/2016	32	32.078	7.75	6.21	49.7	0.846	7.72	531
5/17/2016	34	34.032	6.11	5.55	44.8	0.874	7.70	1383
5/17/2016	36	36.100	6.14	3.93	31.6	0.922	7.57	1227
5/17/2016	38	38.009	5.99	2.83	22.6	0.984	7.49	885
5/17/2016	40	40.200	5.96	2.54	20.1	1.006	7.46	846
5/17/2016	42	42.038	5.96	2.06	16.4	1.013	7.43	359
5/17/2016	44	42.038	6.01	1.74	14.0	1.040	7.39	889

Date	Depth	Dep25	Temp	DO	DO%	SpCond	pH	BG
MMDDYY	feet	feet	°C	mg/l	Sat	mS/cm	Units	
6/14/2016	0	0.500	20.90	9.41	105.6	0.750	8.28	656
6/14/2016	1	1.076	20.90	9.42	105.6	0.750	8.21	743
6/14/2016	2	2.142	20.90	9.41	105.6	0.750	8.18	1425
6/14/2016	3	3.028	20.90	9.42	105.6	0.750	8.17	1167
6/14/2016	4	4.073	20.89	9.42	105.7	0.750	8.17	899
6/14/2016	6	6.048	20.80	9.41	103.0	0.749	8.14	315
6/14/2016	8	8.019	20.71	9.22	103.1	0.750	8.15	582
6/14/2016	10	10.090	20.50	9.21	102.4	0.751	8.15	375
6/14/2016	12	12.022	19.83	9.01	99.2	0.754	8.14	911
6/14/2016	14	14.027	18.34	8.98	95.9	0.763	8.03	228
6/14/2016	16	16.179	15.95	9.08	92.1	0.781	7.97	427
6/14/2016	18	18.096	13.66	9.22	88.4	0.789	7.97	930
6/14/2016	20	20.040	11.43	8.90	80.1	0.795	7.93	967
6/14/2016	22	22.040	9.55	8.32	73.9	0.811	7.88	1545
6/14/2016	24	24.124	8.39	7.78	66.1	0.818	8.82	1097
6/14/2016	26	26.025	7.70	6.40	53.1	0.839	7.65	585
6/14/2016	28	28.125	7.39	5.80	48.2	0.862	7.61	797
6/14/2016	30	30.120	7.20	5.11	42.0	0.895	7.58	1842
6/14/2016	32	32.078	7.05	3.99	32.6	0.957	7.53	801
6/14/2016	34	34.032	6.98	3.16	25.2	1.001	7.48	1650
6/14/2016	36	36.100	6.92	2.42	19.6	1.029	7.45	2347
6/14/2016	38	38.009	6.79	1.91	15.5	1.090	7.40	743
6/14/2016	40	40.200	6.79	1.26	10.2	1.154	7.36	935
6/14/2016	42	42.038	6.66	0.79	6.5	1.193	7.31	1574

Date	Depth	Dep25	Temp	DO	DO%	SpCond	pH	BG
MMDDYY	feet	feet	øC	mg/l	Sat	mS/cm	Units	
7/12/2016	0	0.418	24.40	8.67	104.0	0.757	8.28	868
7/12/2016	1	0.980	24.38	8.67	104.0	0.756	8.28	1032
7/12/2016	2	2.178	24.38	8.68	104.0	0.756	8.28	-1018
7/12/2016	3	3.016	24.39	8.68	104.1	0.757	8.28	797
7/12/2016	4	4.045	24.39	8.70	104.3	0.756	8.28	-2632
7/12/2016	6	6.029	24.38	8.71	104.5	0.756	8.28	-838
7/12/2016	8	7.966	24.37	8.75	104.9	0.756	8.28	-701
7/12/2016	10	10.011	24.33	8.80	105.4	0.756	8.28	2028
7/12/2016	12	12.044	24.23	8.71	104.1	0.755	8.28	-277
7/12/2016	14	13.942	23.59	7.83	92.5	0.761	8.16	-1454
7/12/2016	16	16.004	19.83	6.25	68.7	0.788	7.89	-1176
7/12/2016	18	18.161	16.80	6.32	65.3	0.802	7.83	1001
7/12/2016	20	20.046	14.19	6.59	64.4	0.812	7.76	1173
7/12/2016	22	21.873	11.25	6.07	55.5	0.836	7.71	2481
7/12/2016	24	23.894	9.48	4.51	39.6	0.863	7.64	665
7/12/2016	26	26.046	8.74	3.98	34.3	0.892	7.58	910
7/12/2016	28	27.972	8.32	2.41	20.6	0.930	7.52	3485
7/12/2016	30	29.996	7.92	1.99	16.8	1.005	7.46	1429
7/12/2016	32	31.985	7.60	1.37	11.5	1.069	7.41	2212
7/12/2016	34	34.076	7.46	0.86	7.2	1.103	7.36	1397
7/12/2016	36	36.100	7.34	0.59	5.0	1.151	7.32	2260

Date	Depth	Dep25	Temp	DO	DO%	SpCond	pH	BG
8/16/2016	0	0.370	24.91	7.96	96.4	0.748	8.24	228
8/16/2016	1	1.038	24.90	7.96	96.4	0.748	8.22	-1587
8/16/2016	2	2.019	24.90	7.96	96.3	0.748	8.21	-1048
8/16/2016	3	3.023	24.90	7.96	96.3	0.748	8.20	940
8/16/2016	4	4.004	24.90	7.95	96.2	0.747	8.20	-997
8/16/2016	6	6.009	24.89	7.94	96.1	0.747	8.20	1785
8/16/2016	8	8.088	24.89	7.93	96.0	0.747	8.20	-2136
8/16/2016	10	10.030	24.88	7.92	95.8	0.747	8.20	-18
8/16/2016	12	12.026	24.87	7.89	95.5	0.747	8.20	-3372
8/16/2016	14	14.083	24.81	7.78	94.0	0.747	8.19	-255
8/16/2016	16	16.071	23.35	5.38	63.2	0.769	8.01	-602
8/16/2016	18	18.058	19.23	3.38	36.7	0.807	7.87	-593
8/16/2016	20	20.022	16.29	3.34	34.2	0.828	7.75	1125
8/16/2016	22	22.060	13.46	3.11	29.9	0.855	7.69	-379
8/16/2016	24	24.038	11.59	2.40	22.1	0.888	7.63	128
8/16/2016	26	26.014	10.00	1.42	12.6	0.942	7.58	1600
8/16/2016	28	28.034	9.14	0.89	7.7	0.982	7.53	1364
8/16/2016	30	30.042	8.62	0.65	5.6	1.066	7.44	2970
8/16/2016	32	32.119	8.15	0.49	4.2	1.13	7.39	4335
8/16/2016	34	34.033	7.95	0.50	4.3	1.161	7.37	2842
8/16/2016	36	36.015	7.84	0.50	4.2	1.182	7.34	1698
8/16/2016	38	38.031	7.64	0.46	3.9	1.219	7.32	1022
8/16/2016	40	40.042	7.25	0.39	3.3	1.265	7.29	2124
8/16/2016	42	42.045	7.10	0.34	2.8	1.284	7.27	4496
8/16/2016	44	44.036	6.98	0.34	2.8	1.291	7.23	7621

Date	Depth	Dep25	Temp	DO	DO%	SpCond	pH	BG
9/13/2016	0	0.432	21.53	8.25	93.7	0.732	8.54	-1160
9/13/2016	1	1.065	21.56	8.23	93.5	0.733	8.43	-971
9/13/2016	2	2.020	21.56	8.22	93.4	0.732	8.38	-2313
9/13/2016	3	3.039	21.56	8.22	93.4	0.733	8.35	-1330
9/13/2016	4	4.016	21.56	8.21	93.3	0.733	8.31	-1485
9/13/2016	6	6.031	21.56	8.21	93.3	0.732	8.27	-1766
9/13/2016	8	8.031	21.56	8.20	93.2	0.732	8.25	-741
9/13/2016	10	8.032	21.57	8.19	93.1	0.732	8.24	-375
9/13/2016	12	10.047	21.57	8.19	93.1	0.732	8.24	-1642
9/13/2016	14	12.093	21.56	8.18	93.0	0.732	8.23	-1049
9/13/2016	16	14.010	21.56	8.18	92.9	0.731	8.23	-263
9/13/2016	18	16.016	21.55	8.14	92.5	0.732	8.21	319
9/13/2016	20	18.076	21.49	7.56	85.7	0.732	8.17	439
9/13/2016	22	20.074	18.00	2.25	23.8	0.814	7.85	408
9/13/2016	24	22.056	14.93	1.23	12.2	0.859	7.72	-106
9/13/2016	26	24.124	12.48	0.87	8.2	0.906	7.61	101
9/13/2016	28	26.097	11.01	0.61	5.6	0.954	7.53	3916
9/13/2016	30	28.062	9.86	0.48	4.2	1.018	7.48	1393
9/13/2016	32	30.037	9.18	0.41	3.6	1.096	7.43	4977
9/13/2016	34	29.976	9.14	0.35	3.1	1.097	7.40	5108
9/13/2016	36	32.063	8.59	0.28	2.4	1.156	7.35	810
9/13/2016	38	34.047	8.23	0.23	1.9	1.182	7.31	3719
9/13/2016	40	36.039	7.89	0.20	1.7	1.208	7.27	3784

Independence Grove 2016
RTRM: Relative Thermal Resistance to Mixing
RVG: Relative Viscosity Gradient

Date	Depth	Temp	DO	DO%	RTRM	RVG
5/17/2016	0.50	12.39	10.75	100.42	0	0
5/17/2016	1.08	12.39	10.76	100.51	0	0
5/17/2016	2.14	12.40	10.75	100.65	-1	-9
5/17/2016	3.03	12.40	10.75	100.65	0	0
5/17/2016	4.07	12.40	10.74	100.55	0	0
5/17/2016	6.05	12.39	10.73	100.23	1	9
5/17/2016	8.02	12.39	10.72	100.14	0	0
5/17/2016	10.09	12.34	10.70	99.95	0	0
5/17/2016	12.02	12.65	10.62	99.88	-4	-26
5/17/2016	14.03	12.29	10.52	98.05	6	34
5/17/2016	16.18	12.21	10.50	97.87	0	0
5/17/2016	18.10	11.86	10.44	96.43	6	34
5/17/2016	20.04	11.18	10.05	91.35	9	60
5/17/2016	22.04	7.42	9.90	82.36	35	325
5/17/2016	24.12	6.66	8.75	71.37	5	80
5/17/2016	26.02	6.44	8.23	66.80	1	20
5/17/2016	28.13	6.24	7.81	63.09	1	20
5/17/2016	30.12	6.19	7.35	59.23	0	10
5/17/2016	32.08	7.75	6.21	52.05	-9	-160
5/17/2016	34.03	6.11	5.55	44.72	9	160
5/17/2016	36.10	6.14	3.93	31.67	0	0
5/17/2016	38.01	5.99	2.83	22.69	1	20
5/17/2016	40.20	5.96	2.54	20.36	0	0
5/17/2016	42.04	5.96	2.06	16.52	0	0
5/17/2016	42.04	6.01	1.74	13.99	0	-10

Date	Depth	Temp	DO	DO%	RTRM	RVG
6/14/2016	0.50	20.90	9.41	105.34	0	0
6/14/2016	1.08	20.90	9.42	105.46	0	0
6/14/2016	2.14	20.90	9.41	105.34	0	0
6/14/2016	3.03	20.90	9.42	105.46	0	0
6/14/2016	4.07	20.89	9.42	105.25	3	4
6/14/2016	6.05	20.80	9.41	105.13	0	0
6/14/2016	8.02	20.71	9.22	102.81	3	4
6/14/2016	10.09	20.50	9.21	102.29	5	8
6/14/2016	12.02	19.83	9.01	98.70	18	28
6/14/2016	14.03	18.34	8.98	95.43	36	60
6/14/2016	16.18	15.95	9.08	91.80	52	96
6/14/2016	18.10	13.66	9.22	88.67	42	157
6/14/2016	20.04	11.43	8.90	81.45	33	188
6/14/2016	22.04	9.55	8.32	72.83	22	163
6/14/2016	24.12	8.39	7.78	66.17	11	103
6/14/2016	26.02	7.70	6.40	53.64	4	56
6/14/2016	28.13	7.39	5.80	48.13	3	40
6/14/2016	30.12	7.20	5.11	42.30	1	10
6/14/2016	32.08	7.05	3.99	32.87	1	20
6/14/2016	34.03	6.98	3.16	25.97	0	10
6/14/2016	36.10	6.92	2.42	19.89	0	0
6/14/2016	38.01	6.79	1.91	15.62	1	20
6/14/2016	40.20	6.79	1.26	10.30	0	0
6/14/2016	42.04	6.66	0.79	6.44	0	10

Date	Depth	Temp	DO	DO%	RTRM	RVG
7/12/2016	0.42	24.40	8.67	103.57	0	0
7/12/2016	0.98	24.38	8.67	103.57	0	0
7/12/2016	2.18	24.38	8.68	103.69	0	0
7/12/2016	3.02	24.39	8.68	103.69	0	0
7/12/2016	4.04	24.39	8.70	103.92	0	0
7/12/2016	6.03	24.38	8.71	104.04	0	0
7/12/2016	7.97	24.37	8.75	104.52	0	0
7/12/2016	10.01	24.33	8.80	105.12	0	0
7/12/2016	12.04	24.23	8.71	103.85	3	4
7/12/2016	13.94	23.59	7.83	92.14	21	28
7/12/2016	16.00	19.83	6.25	68.46	102	148
7/12/2016	18.16	16.80	6.32	64.98	72	124
7/12/2016	20.05	14.19	6.59	64.08	50	146
7/12/2016	21.87	11.25	6.07	55.30	44	248
7/12/2016	23.89	9.48	4.51	39.39	20	154
7/12/2016	26.05	8.74	3.98	34.18	7	60
7/12/2016	27.97	8.32	2.41	20.50	3	34
7/12/2016	30.00	7.92	1.99	16.76	3	36
7/12/2016	31.98	7.60	1.37	11.43	3	40
7/12/2016	34.08	7.46	0.86	7.15	1	10
7/12/2016	36.10	7.34	0.59	4.90	1	10

Date	Depth	Temp	DO	DO%	RTRM	RVG
8/16/2016	0.37	24.91	7.96	96.15	1/0/1900	0
8/16/2016	1.04	24.90	7.96	95.97	3	4
8/16/2016	2.02	24.90	7.96	95.97	0	0
8/16/2016	3.02	24.90	7.96	95.97	0	0
8/16/2016	4.00	24.90	7.95	95.85	0	0
8/16/2016	6.01	24.89	7.94	95.73	0	0
8/16/2016	8.09	24.89	7.93	95.61	0	0
8/16/2016	10.03	24.88	7.92	95.49	0	0
8/16/2016	12.03	24.87	7.89	95.13	0	0
8/16/2016	14.08	24.81	7.78	93.80	0	0
8/16/2016	16.07	23.35	5.38	63.07	46	60
8/16/2016	18.06	19.23	3.38	36.58	111	164
8/16/2016	20.02	16.29	3.34	33.98	67	120
8/16/2016	22.06	13.46	3.11	29.78	51	186
8/16/2016	24.04	11.59	2.40	22.01	28	163
8/16/2016	26.01	10.00	1.42	12.58	18	128
8/16/2016	28.03	9.14	0.89	7.72	9	77
8/16/2016	30.04	8.62	0.65	5.57	4	43
8/16/2016	32.12	8.15	0.49	4.15	4	43
8/16/2016	34.03	7.95	0.50	4.21	1	19
8/16/2016	36.01	7.84	0.50	4.20	1	10
8/16/2016	38.03	7.64	0.46	3.85	1	20
8/16/2016	40.04	7.25	0.39	3.23	3	40
8/16/2016	42.04	7.10	0.34	2.80	1	20
8/16/2016	44.04	6.98	0.34	2.79	0	10

Date	Depth	Temp	DO	DO%	RTRM	RVG
9/13/2016	0.43	21.53	8.25	93.44	0	0
9/13/2016	1.06	21.56	8.23	93.22	0	0
9/13/2016	2.02	21.56	8.22	93.10	0	0
9/13/2016	3.04	21.56	8.22	93.10	0	0
9/13/2016	4.02	21.56	8.21	92.99	0	0
9/13/2016	6.03	21.56	8.21	92.99	0	0
9/13/2016	8.03	21.56	8.20	92.88	0	0
9/13/2016	8.03	21.57	8.19	92.76	0	0
9/13/2016	10.05	21.57	8.19	92.76	0	0
9/13/2016	12.09	21.56	8.18	92.65	0	0
9/13/2016	14.01	21.56	8.18	92.65	0	0
9/13/2016	16.02	21.55	8.14	92.20	0	0
9/13/2016	18.08	21.49	7.56	85.46	3	4
9/13/2016	20.07	18.00	2.25	23.77	85	136
9/13/2016	22.06	14.93	1.23	12.17	64	129
9/13/2016	24.12	12.48	0.87	8.15	41	214
9/13/2016	26.10	11.01	0.61	5.53	19	120
9/13/2016	28.06	9.86	0.48	4.23	14	103
9/13/2016	30.04	9.18	0.41	3.56	7	60
9/13/2016	29.98	9.14	0.35	3.04	0	0
9/13/2016	32.06	8.59	0.28	2.39	5	51
9/13/2016	34.05	8.23	0.23	1.95	2	26
9/13/2016	36.04	7.89	0.20	1.68	3	37

Independence Grove IEPA Ranking

TROPHIC STATUS

Carlson's TSIp 40.4 Mesotrophic

IMPAIRMENT ASSESSMENTS

Total Phosphorus None
 Total Nitrogen None
 pH None
 Low DO None
 Total Dissolved Solids None
 Total Suspended Solids None
 Aquatic Plants-Native Yes
 Non-Native Aquatic Plants Yes
 Non-Native Animals Yes

AQUATIC LIFE USE IMPAIRMENT INDEX

Mean Trophic State
 Macrophyte Impairment
 Sediment Impairment (NVSS)
 Degree of Use Support

Weighting Criteria	Points	Overall Use Support Points	Degree of Support
40.4	40		
Substantial	15		
Minimal	0		
	<u>55</u>	0	Full

RECREATION USE IMPAIRMENT INDEX

Mean Trophic State Index
 Macrophyte Impairment
 Sediment Impairment (NVSS)
 Degree of Use Support

40.4	40.4		
Substantial	15		
Minimal	0		
	<u>55.4</u>	0	Full
<u>Overall Use Index</u>		0.00	Full

**Aquatic Plants found at the 147 sampling sites on Independence Grove in July of 2016.
The maximum depth that plants were found was 15 feet.**

Plant Density	American Pondweed	Brittle Naiad	Chara	Coontail	Curlyleaf Pondweed	Duckweed	Elodea	Eurasian Watermilfoil	Floatingleaf Pondweed	Illinois Pondweed
Absent	113	145	135	82	141	146	113	118	146	132
Present	10	2	2	14	6	0	11	10	0	3
Common	22	0	7	28	0	1	15	16	0	5
Abundant	2	0	2	13	0	0	8	2	1	6
Dominant	0	0	1	10	0	0	0	1	0	1
% Plant Occurrence	23.1	1.4	8.2	44.2	4.1	0.7	23.1	19.7	0.7	10.2

Plant Density	Northern Naiad	Sago Pondweed	Slender Naiad	Small Pondweed	Spatterdock	Southern Naiad	White Water Crowfoot	Watermeal	Water Stargrass	White Water Lily
Absent	142	135	139	137	146	145	122	144	117	130
Present	3	4	3	7	0	1	11	0	8	5
Common	1	8	5	3	1	0	8	2	18	4
Abundant	1	0	0	0	0	1	5	1	4	8
Dominant	0	0	0	0	0	0	1	0	0	0
% Plant Occurrence	3.4	8.2	5.4	6.8	0.7	1.4	17.0	2.0	20.4	11.6

Distribution of rake density across all sampling sites.

Rake Density (coverage)	# of Sites	% of Sites
No Plants	41	28
>0-10%	8	5
10-40%	27	18
40-60%	31	21
60-90%	31	21
>90%	9	6
Total Sites with Plants	106	72
Total # of Sites	147	100

Independence Grove Lake 2016 Land Use

LanduseName	ACRES
Agricultural	20.0
Forest and Grassland	54.0
Government and Institutional	5.1
Public and Private Open Space	113.2
Single Family	0.2
Transportation	15.6
Water	126.7

Land Use	Acreage	% of Total
Agricultural	20.0	6.0%
Forest and Grassland	54.0	16.1%
Government and Institutional	5.1	1.5%
Public and Private Open Space	113.2	33.8%
Single Family	0.2	0.1%
Transportation	15.6	4.7%
Water	126.7	37.9%
Total Acres	334.79	100.0%

Land Use	Acreage	Runoff Coeff.	Estimated Runoff, acft.	% Total of Estimated Runoff
Agricultural	19.97	0.05	2.7	0.6
Forest and Grassland	53.96	0.50	74.2	16.9
Government and Institutional	5.06	0.50	7.0	1.6
Public and Private Open Space	113.22	0.15	46.7	10.6
Single Family	0.22	0.30	0.2	0.0
Transportation	15.63	0.30	12.9	2.9
Water	126.74	0.85	296.3	67.3
TOTAL	334.79		439.9	100.0

Lake volume

1945.00 acre-feet

Retention Time (years)= lake volume/runoff

4.42 years

1613.75 days

Table 1: Independence Grove Lake Shoreline Erosion Condition 2016

Reach	None		Slight		Moderate		Severe		Total	Lateral Recession Rate
	Linear ft.	% Reach	Linear Ft.	% Reach		% Reach	Linear Ft.	% Reach		
IDG01	94.8	3%	1393.0	50%	689.5	25%	591.8	21%	2769.1	0.12
IDG02	465.2	53%	306.8	35%	110.5	13%	0.0	0%	882.5	0.03
IDG03	14.0	1%	1604.1	88%	203.4	11%	0.0	0%	1821.5	0.03
IDG04	1076.4	91%	104.4	9%	0.0	0%	0.0	0%	1180.8	0.01
IDG05	532.9	33%	1029.5	64%	48.6	3%	0.0	0%	1611.0	0.02
IDG06	72.5	11%	472.8	73%	100.7	16%	0.0	0%	646.0	0.03
IDG07	706.3	97%	18.6	3%	0.0	0%	0.0	0%	724.9	0.01
IDG08	1082.8	64%	389.3	23%	210.3	13%	0.0	0%	1682.4	0.02
IDG09	637.7	50%	606.1	48%	22.8	2%	0.0	0%	1266.6	0.02
IDG10	606.6	39%	788.2	51%	142.8	9%	0.0	0%	1537.6	0.03
IDG11	275.2	54%	234.1	46%	0.0	0%	0.0	0%	509.3	0.02
IDG12	572.0	77%	100.3	14%	68.3	9%	0.0	0%	740.6	0.02
IDG13	490.0	61%	314.4	39%	0.0	0%	0.0	0%	804.4	0.02
IDG14	656.0	83%	0.0	0%	131.0	17%	0.0	0%	787.0	0.02
IDG15	449.9	49%	363.6	39%	107.7	12%	0.0	0%	921.2	0.03
IDG16	534.0	62%	236.9	28%	88.8	10%	0.0	0%	859.7	0.02
IDG17	475.9	41%	426.7	36%	269.6	23%	0.0	0%	1172.2	0.03
IDG18	147.6	12%	858.4	69%	241.2	19%	0.0	0%	1247.2	0.04
IDG19	92.2	22%	323.1	78%	0.0	0%	0.0	0%	415.3	0.02
IDG20	557.2	60%	367.5	40%	0.0	0%	0.0	0%	924.7	0.02
IDG21	1250.5	74%	437.5	26%	0.0	0%	0.0	0%	1688.0	0.01
Total	10789.7	45%	10375.3	43%	2435.2	10%	591.8	2%	24192.0	0.03

Table 2: Independence Grove Lake Shoreline Buffer Condition 2016

Reach Code	Good Condition (ft/%)		Fair Condition (ft/%)		Poor Condition (ft/%)		Shoreline Length Assessed (ft)
	ft	%	ft	%	ft	%	
IDG01	375.2	67.5	164.7	29.6	16.0	2.9	555.9
IDG02	58.8	11.7	443.0	88.3	0.0	0.0	501.8
IDG03	472.3	100.0	0.0	0.0	0.0	0.0	472.3
IDG04	405.7	81.5	45.4	9.1	46.5	9.3	497.6
IDG05	399.0	81.3	83.9	17.1	8.0	1.6	490.9
IDG06	45.3	23.0	0.0	0.0	151.6	77.0	196.9
IDG07	0.0	0.0	167.5	75.8	53.4	24.2	220.9
IDG08	184.7	35.9	274.2	53.2	56.1	10.9	515.0
IDG09	160.5	41.8	122.3	31.9	100.8	26.3	383.6
IDG10	261.5	55.8	207.0	44.2	0.0	0.0	468.5
IDG11	138.5	89.2	16.7	10.8	0.0	0.0	155.2
IDG12	51.1	22.7	61.1	27.1	113.4	50.3	225.6
IDG13	51.4	21.0	193.7	79.0	0.0	0.0	245.1
IDG14	244.6	100.0	0.0	0.0	0.0	0.0	244.6
IDG15	195.2	70.8	80.7	29.2	0.0	0.0	275.9
IDG16	236.3	90.2	18.9	7.2	6.8	2.6	262.0
IDG17	207.9	58.2	142.1	39.8	7.2	2.0	357.2
IDG18	277.0	72.9	76.3	20.1	26.8	7.1	380.1
IDG19	27.4	21.6	30.3	23.9	68.9	54.4	126.6
IDG20	250.5	88.9	31.3	11.1	0.0	0.0	281.8
IDG21	476.5	92.6	37.9	7.4	0.0	0.0	514.4

Appendix C:
Methods for Field Data Collection and Laboratory Analyses

Water Sampling and Laboratory Analyses

Two water samples were collected once a month from May through September. Sample locations were at the deepest point in the lake (see sample site map), three feet below the surface, and 3 feet above the bottom. Samples were collected with a horizontal Van Dorn water sampler. Approximately three liters of water were collected for each sample for all lab analyses. After collection, all samples were placed in a cooler with ice until delivered to the Lake County Health Department lab, where they were refrigerated. Analytical methods for the parameters are listed in Table A1. Except nitrate nitrogen, all methods are from the Eighteenth Edition of Standard Methods, (eds. American Public Health Association, American Water Works Association, and Water Pollution Control Federation, 1992). Methodology for nitrate nitrogen was taken from the 14th edition of Standard Methods. Dissolved oxygen, temperature, conductivity and pH were measured at the deep hole with a Hydrolab DataSonde® 4a. Photosynthetic Active Radiation (PAR) was recorded using a LI-COR® 192 Spherical Sensor attached to the Hydrolab DataSonde® 4a. Readings were taken at the surface and then every two feet until reaching the bottom.

Plant Sampling

In order to randomly sample each lake, mapping software (ArcMap 9.3) overlaid a grid pattern onto an aerial photo of Lake County and placed points 60 or 30 meters apart, depending on lake size. Plants were sampled using a garden rake fitted with hardware cloth. The hardware cloth surrounded the rake tines and is tapered two feet up the handle. A rope was tied to the end of the handle for retrieval. At designated sampling sites, the rake was tossed into the water, and using the attached rope, was dragged across the bottom, toward the boat. After pulling the rake into the boat, plant coverage was assessed for overall abundance. Then plants were individually identified and placed in categories based on coverage. Plants that were not found on the rake but were seen in the immediate vicinity of the boat at the time of sampling were also recorded. Plants difficult to identify in the field were placed in plastic bags and identified with plant keys after returning to the office. The depth of each sampling location was measured either by a hand-held depth meter, or by pushing the rake straight down and measuring the depth along the rope or rake handle. One-foot increments were marked along the rope and rake handle to aid in depth estimation.

Plankton Sampling

Plankton were sampled at the same location as water quality samples. Using the Hydrolab DataSonde® 4a or YSI 6600 Sonde® 1% light level depth (depth where the water light is 1% of the surface irradiance) was determined. A plankton net/tow, with 63µm mesh, was then lowered to the pre-determined 1% light level depth and retrieved vertically. On the way up the water column, plankton were collected within a small cup on the bottom of the tow. The collected sample was then emptied into a pre-labeled brown plastic bottle. The net was rinsed with deionized water into the bottle in order to ensure all the plankton were collected. The sample was then transferred to a graduated cylinder to measure the amount of milliliters (mL) that the sample was. The sample was then returned to the bottle and preserved with Lugol's iodine solution (5 drops/mL). The sample bottle was then closed and stored in a cooler until returning

to the lab, where it was transferred to the refrigerator until enumeration. Enumeration was performed within three months, but ideally within one month, under a microscope. Prior to sub-sample being removed for enumeration, the sample bottle was inverted several times to ensure proper homogenization. An automated pipette was used to retrieve 1 mL of sample, which was then placed in a Sedgewick Rafter slide. This is a microscope slide on which a rectangular chamber has been constructed, measuring 50 mm x 20 mm in area and 1 mm deep. The slide was then placed under the microscope and counted at a 20X magnification (phytoplankton) or 10X magnification (zooplankton). For phytoplankton, twenty fields of view were randomly counted with all species within each field counted. Due to their larger size, zooplankton were counted throughout the entire slide. Through calculations, it was determined how many of each species were in 1 mL of lake water.

Phytoplankton (algae) are free-floating and microscopic and are distinguished from plants because they lack roots, stems and leaves. There are four distinct groups of phytoplankton found in Lake County lakes: blue-greens, greens, diatoms, and dinoflagellates/chrysophytes. Blue-greens are also known as cyanobacteria because they are the only group of bacteria that obtain their energy from photosynthesis like plants. Some of these species can be toxic. Green algae are the closest ancestors of land plants and are the most common group. Diatoms are unique because they are encased in a cell wall made of silica that can be very ornate. Dinoflagellates and chrysophytes are almost always flagellated (able to move by flagella, a whip-like tail) and some can both photosynthesize and consume bacteria for food.

Zooplankton are made up of rotifers and two crustacean groups; the cladocerans and the copepods (broken down further into calanoids and cyclopoids). Rotifers are smaller and most have a crown of cilia (hair-like structure) used for movement and drawing in suspended particles to eat. Crustaceans have jointed appendages and are enclosed in an exoskeleton. Cladocerans, such as the “water flea” *Daphnia* species, are filter-feeding like rotifers, while the copepod group contains both filter-feeders (calanoids and cyclopoids) and raptorial species (cyclopoids).

Shoreline Assessment

Shoreline Assessment Protocol

Each lake was divided into reaches in ArcGIS based on nearshore landuse. For each reach, a shoreline assessment worksheet was filled out in the field focusing on shoreline conditions (land use, slope, erosion, buffer, etc) that describe the overall reach segment of the lake.

A GPS Trimble unit was used to collect the degree of shoreline erosion along the entire length of the lake. The degree of shoreline erosion was categorically defined as none, slight, moderate, or severe. Below are brief descriptions of each category.

Table 1: Degree of Shoreline Erosion

Category	Description
None	Includes man-made erosion control such as rip-rap and sea wall.
Slight	Minimal or no observable erosion; generally considered stable; no erosion control practices will be recommended with the possible exception of small problem areas noted within an area otherwise designated as "slight". Beaches have been included as "slight" erosion.
Moderate	Recession is characterized by past or recently eroded banks; area may exhibit some exposed roots, fallen vegetation or minor slumping of soil material; erosion control practices may be recommended although the section is not deemed to warrant immediate remedial action.
Severe	Recession is characterized by eroding of exposed soil on nearly vertical banks, exposed roots, fallen vegetation or extensive slumping of bank material, undercutting, washouts or fence posts exhibiting realignment; erosion control practices are recommended and immediate remedial action may be warranted.

Lateral recession rates were calculated on a per reach basis based on the IL EPA stream methodology, defining lateral recession into four main categories (slight, moderate, severe, and very severe). Descriptions of each category are defined in the Table 2.

Table 2: Lateral Recession Rate Categories

Lateral Recession Rate	Description	Description
0.01 – 0.05	Slight	Some bare bank but active erosion not readily apparent. Some rills but no vegetation overhanging. No exposed tree roots.
0.06 – 0.2	Moderate	Bank mostly bare with some rills and vegetation overhanging.
0.3 – 0.5	Severe	Bank is bare with rills and severe vegetative overhang. Many exposed tree roots and some fallen trees and slumps or slips. Some changes in cultural features such as fence corners missing and realignment of roads or trails. Channel cross-section becomes more U-shaped as opposed to V-shaped.
0.5+	Very Severe	Bank is bare with gullies and severe vegetative overhang. Many fallen trees, drains and culverts eroding out and changes in cultural features as above. Massive slips or washouts common. Channel cross-section is U-shaped and streamcourse or gully may be meandering.

Shoreline Buffer Condition

Lakeshore buffer condition was assessed using a qualitative methodology that considered an area up to 25 feet inland from the shoreline for each reach. The assessment was done by viewing high resolution 2014 aerial images in ArcGIS. A 25 foot buffer was chosen based on research that indicates a 25-foot vegetated buffer is the minimum effective width for in-lake habitat maintenance (a 15 foot buffer is the minimum effective width for bank stability). Criteria used for category assignment are shown in table below.

Table 3: Shoreline Buffer Condition Categories

Category	Criteria	Percentage
Good	Unmowed grasses & forbs + tree trunks + shrubs <i>and</i> impervious surfaces	$\geq 70\%$
		$\leq 5\%$
Fair	Unmowed grasses & forbs + tree trunks + shrubs <i>and</i> Impervious surface	$\geq 50\%$ and $< 70\%$
		$\leq 10\%$
Poor	Unmowed grasses & forbs + tree trunks + shrubs <i>and</i> Impervious surface	$< 50\%$
		$\geq 50\%$

Wildlife Assessment

Species of wildlife were noted during visits to each lake. When possible, wildlife was identified to species by sight or sound. However, due to time constraints, collection of quantitative information was not possible. Thus, all data should be considered anecdotal. Some of the species on the list may have only been seen once, or were spotted during their migration through the area.

Table A1. Analytical methods used for water quality parameters.

<i>Parameter</i>	<i>Method</i>
Temperature	Hydrolab DataSonde® 4a or YSI 6600 Sonde®
Dissolved oxygen	Hydrolab DataSonde® 4a or YSI 6600 Sonde®
Nitrate and Nitrite nitrogen	USEPA 353.2 rev. 2.0 EPA-600/R-93/100 Detection Limit = 0.05 mg/L
Ammonia nitrogen	SM 18 th ed. Electrode method, #4500 NH ₃ -F Detection Limit = 0.1 mg/L
Total Kjeldahl nitrogen	SM 18 th ed, 4500-N _{org} C Semi-Micro Kjeldahl, plus 4500 NH ₃ -F Detection Limit = 0.5 mg/L
pH	Hydrolab DataSonde® 4a, or YSI 6600 Sonde® Electrometric method
Total solids	SM 18 th ed, Method #2540B
Total suspended solids	SM 18 th ed, Method #2540D Detection Limit = 0.5 mg/L
Chloride	SM 18 th ed, Method #4500C1-D
Total volatile solids	SM 18 th ed, Method #2540E, from total solids
Alkalinity	SM 18 th ed, Method #2320B, potentiometric titration curve method
Conductivity	Hydrolab DataSonde® 4a or YSI 6600 Sonde®
Total phosphorus	SM 18 th ed, Methods #4500-P B 5 and #4500-P E Detection Limit = 0.01 mg/L
Soluble reactive phosphorus	SM 18 th ed, Methods #4500-P B 1 and #4500-P E Detection Limit = 0.005 mg/L
Clarity	Secchi disk
Color	Illinois EPA Volunteer Lake Monitoring Color Chart
Photosynthetic Active Radiation (PAR)	Hydrolab DataSonde® 4a or YSI 6600 Sonde®, LI-COR® 192 Spherical Sensor

**APPENDIX D. INTERPRETING YOUR LAKE'S WATER QUALITY
DATA**

Lakes possess a unique set of physical and chemical characteristics that will change over time. These in-lake water quality characteristics, or parameters, are used to describe and measure the quality of lakes, and they relate to one another in very distinct ways. As a result, it is virtually impossible to change any one component in or around a lake without affecting several other components, and it is important to understand how these components are linked.

The following pages will discuss the different water quality parameters measured by Lake County Health Department staff, how these parameters relate to each other, and why the measurement of each parameter is important. The median values (the middle number of the data set, where half of the numbers have greater values, and half have lesser values) of data collected from Lake County lakes from 2000-2010 will be used in the following discussion.

Temperature and Dissolved Oxygen:

Water temperature fluctuations will occur in response to changes in air temperatures, and can have dramatic impacts on several parameters in the lake. In the spring and fall, lakes tend to have uniform, well-mixed conditions throughout the water column (surface to the lake bottom). However, during the summer, deeper lakes will separate into distinct water layers. As surface water temperatures increase with increasing air temperatures, a large density difference will form between the heated surface water and colder bottom water. Once this difference is large enough, these two water layers will separate and generally will not mix again until the fall. At this time the lake is thermally stratified. The warm upper water layer is called the *epilimnion*, while the cold bottom water layer is called the *hypolimnion*. In some shallow lakes, stratification and destratification can occur several times during the summer. If this occurs the lake is described as polymictic. Thermal stratification also occurs to a lesser extent during the winter, when warmer bottom water becomes separated from ice-forming water at the surface until mixing occurs during spring ice-out.

Monthly temperature profiles were established on each lake by measuring water temperature every foot (lakes \leq 15 feet deep) or every two feet (lakes $>$ 15 feet deep) from the lake surface to the lake bottom. These profiles are important in understanding the distribution of chemical/biological characteristics and because increasing water temperature and the establishment of thermal stratification have a direct impact on dissolved oxygen (DO) concentrations in the water column. If a lake is shallow and easily mixed by wind, the DO concentration is usually consistent throughout the water column. However, shallow lakes are typically dominated by either plants or algae, and increasing water temperatures during the summer speeds up the rates of photosynthesis and decomposition in surface waters. Many of the plants or algae die at the end of the growing season. Their decomposition results in heavy oxygen consumption and can lead to an oxygen crash. In deeper, thermally stratified lakes, oxygen production is greatest in the top portion of the lake, where sunlight drives photosynthesis, and oxygen consumption is greatest near the bottom of a lake, where sunken organic matter accumulates and decomposes. The oxygen difference between the top and bottom water layers can be dramatic, with plenty of oxygen near the surface, but practically none near the bottom.

The oxygen profiles measured during the water quality study can illustrate if this is occurring. This is important because the absence of oxygen (anoxia) near the lake bottom can have adverse effects in eutrophic lakes resulting in the chemical release of phosphorus from lake sediment and the production of hydrogen sulfide (rotten egg smell) and other gases in the bottom waters. Low oxygen conditions in the upper water of a lake can also be problematic since all aquatic organisms need oxygen to live. Some oxygen may be present in the water, but at too low a concentration to sustain aquatic life. Oxygen is needed by all plants, virtually all algae and for many chemical reactions that are important in lake functioning. Most adult sport-fish such as largemouth bass and bluegill require at least 3 mg/L of DO in the water to survive. However, their offspring require at least 5 mg/L DO as they are more sensitive to DO stress. When DO concentrations drop below 3 mg/L, rough fish such as carp and green sunfish are favored and over time will become the dominant fish species.

External pollution in the form of oxygen-demanding organic matter (i.e., sewage, lawn clippings, soil from shoreline erosion, and agricultural runoff) or nutrients that stimulate the growth of excessive organic matter (i.e., algae and plants) can reduce average DO concentrations in the lake by increasing oxygen consumption. This can have a detrimental impact on the fish community, which may be squeezed into a very small volume of water as a result of high temperatures in the epilimnion and low DO levels in the hypolimnion.

Nutrients:

Phosphorus:

For most Lake County lakes, phosphorus is the nutrient that limits plant and algae growth. This means that any addition of phosphorus to a lake will typically result in algae blooms or high plant densities during the summer. The source of phosphorus to a lake can be external or internal (or both). External sources of phosphorus enter a lake through point (i.e., storm pipes and wastewater discharge) and non-point runoff (i.e., overland water flow). This runoff can pick up large amounts of phosphorus from agricultural fields, septic systems or impervious surfaces before it empties into the lake.

Internal sources of phosphorus originate within the lake and are typically linked to the lake sediment. In lakes with high oxygen levels (oxic), phosphorus can be released from the sediment through plants or sediment resuspension. Plants take up sediment-bound phosphorus through their roots, releasing it in small amounts to the water column throughout their life cycles, and in large amounts once they die and begin to decompose. Sediment resuspension can occur through biological or mechanical means. Bottom-feeding fish, such as common carp and black bullhead can release phosphorus by stirring up bottom sediment during feeding activities and can add phosphorus to a lake through their fecal matter. Sediment resuspension, and subsequent phosphorus release, can also occur via wind/wave action or through the use of artificial aerators, especially in shallow lakes. In lakes that thermally stratify, internal phosphorus release can occur from the sediment through chemical means. Once oxygen is depleted (anoxia) in the hypolimnion, chemical reactions occur in which phosphorus bound to iron complexes in the

sediment becomes soluble and is released into the water column. This phosphorus is trapped in the hypolimnion and is unavailable to algae until fall turnover, and can cause algae blooms once it moves into the sunlit surface water at that time. Accordingly, many of the lakes in Lake County are plagued by dense algae blooms and excessive, exotic plant coverage, which negatively affect DO levels, fish communities and water clarity.

Lakes with an average phosphorus concentration greater than 0.05 mg/L are considered nutrient rich. The median near surface total phosphorus (TP) concentration in Lake County lakes from 2000-2010 was 0.065 mg/L and ranged from a non-detectable minimum of <0.010 mg/L on seven lakes to a maximum of 3.880 mg/L on Albert Lake. The median anoxic TP concentration in Lake County lakes from 2000-2010 was 0.174 mg/L and ranged from a minimum of 0.012 mg/L in Independence Grove Lake to a maximum of 3.800 mg/L in Taylor Lake.

The analysis of phosphorus also included soluble reactive phosphorus (SRP), a dissolved form of phosphorus that is readily available for plant and algae growth. SRP is not discussed in great detail in most of the water quality reports because SRP concentrations vary throughout the season depending on how plants and algae absorb and release it. It gives an indication of how much phosphorus is available for uptake, but, because it does not take all forms of phosphorus into account, it does not indicate how much phosphorus is truly present in the water column. TP is considered a better indicator of a lake's nutrient status because its concentrations remain more stable than soluble reactive phosphorus. However, elevated SRP levels are a strong indicator of nutrient problems in a lake.

Nitrogen:

Nitrogen is also an important nutrient for plant and algae growth. Sources of nitrogen to a lake vary widely, ranging from fertilizer and animal wastes, to human waste from sewage treatment plants or failing septic systems, to groundwater, air and rainfall. As a result, it is very difficult to control or reduce nitrogen inputs to a lake. Different forms of nitrogen are present in a lake under different oxic conditions. NH_4^+ (ammonium) is released from decomposing organic material under anoxic conditions and accumulates in the hypolimnion of thermally stratified lakes. If NH_4^+ comes into contact with oxygen, it is immediately converted to NO_2^- (nitrite) which is then oxidized to NO_3^- (nitrate). Therefore, in a thermally stratified lake, levels of NH_4^+ would only be elevated in the hypolimnion and levels of NO_3^- would only be elevated in the epilimnion. Both NH_4^+ and NO_3^- can be used as a nitrogen source by aquatic plants and algae. Total Kjeldahl nitrogen (TKN) is a measure of organic nitrogen plus ammonium. Adding the concentrations of TKN and nitrate together gives an indication of the amount of total nitrogen present in the water column. If inorganic nitrogen (NO_3^- , NO_2^- , NH_4^+) concentrations exceed 0.3 mg/L in spring, sufficient nitrogen is available to support summer algae blooms. However, low nitrogen levels do not guarantee limited algae growth the way low phosphorus levels do. Nitrogen gas in the air can dissolve in lake water and blue-green algae can "fix" atmospheric nitrogen, converting it into a usable form. Since other types of algae do not have the ability to do this, nuisance blue-green algae blooms are typically associated with lakes that are nitrogen limited (i.e., have low nitrogen levels).

The ratio of TKN plus nitrate nitrogen to total phosphorus (TN:TP) can indicate whether plant/algae growth in a lake is limited by nitrogen or phosphorus. Ratios of less than 10:1 suggest a system limited by nitrogen, while lakes with ratios greater than 20:1 are limited by phosphorus. It is important to know if a lake is limited by nitrogen or phosphorus because any addition of the limiting nutrient to the lake will, likely, result in algae blooms or an increase in plant density.

Solids:

Although several forms of solids (total solids, total suspended solids, total volatile solids, total dissolved solids) were measured each month by the Lakes Management Staff, total suspended solids (TSS) and total volatile solids (TVS) have the most impact on other variables and on the lake as a whole. TSS are particles of algae or sediment suspended in the water column. High TSS concentrations can result from algae blooms, sediment resuspension, and/or the inflow of turbid water, and are typically associated with low water clarity and high phosphorus concentrations in many lakes in Lake County. Low water clarity and high phosphorus concentrations, in turn, exacerbate the high TSS problem by leading to reduced plant density (which stabilize lake sediment) and increased occurrence of algae blooms. The median TSS value in epilimnetic waters in Lake County was 8.1 mg/L, ranging from below the 0.1 mg/L detection limit to 165 mg/L in Fairfield Marsh.

TVS represents the fraction of total solids that are organic in nature, such as algae cells, tiny pieces of plant material, and/or tiny animals (zooplankton) in the water column. High TVS values indicate that a large portion of the suspended solids may be made up of algae cells. This is important in determining possible sources of phosphorus to a lake. If much of the suspended material in the water column is determined to be resuspended sediment that is releasing phosphorus, this problem would be addressed differently than if the suspended material was made up of algae cells that were releasing phosphorus. The median TVS value was 123.0 mg/L, ranging from 34.0 mg/L in Pulaski Pond to 298.0 mg/L in Fairfield Marsh.

Total dissolved solids (TDS) are the amount of dissolved substances, such as salts or minerals, remaining in water after evaporation. These dissolved solids are discussed in further detail in the *Alkalinity* and *Conductivity* sections of this document. TDS concentrations were measured in Lake County lakes prior to 2004. This practice was discontinued due to the strong correlation of TDS to conductivity and chloride concentrations. Since 2004, chloride concentrations data are collected..

Water Clarity:

Water clarity (transparency) is not a chemical property of lake water, but is often an indicator of a lake's overall water quality. It is affected by a lake's water color, which is a reflection of the amount of total suspended solids and dissolved organic chemicals. Thus, transparency is a

measure of particle concentration and is measured with a Secchi disk. Generally, the lower the clarity or Secchi depth, the poorer the water quality. A decrease in Secchi depth during the summer occurs as the result of an increase in suspended solids (algae or sediment) in the water column. Aquatic plants play an important role in the level of water clarity and can, in turn, be negatively affected by low clarity levels. Plants increase clarity by competing with algae for resources and by stabilizing sediments to prevent sediment resuspension. A lake with a healthy plant community will almost always have higher water clarity than a lake without plants. Additionally, if the plants in a lake are removed (through herbicide treatment or the stocking of grass carp), the lake will probably become dominated by algae and Secchi depth will decrease. This makes it very difficult for plants to become re-established due to the lack of available sunlight and the lake will, most likely, remain turbid. Turbidity will be accelerated if the lake is very shallow and/or common carp are present. Shallow lakes are more susceptible to sediment resuspension through wind/wave action and are more likely to experience clarity problems if plants are not present to stabilize bottom sediment.

Common Carp are prolific fish that feed on invertebrates in the sediment. Their feeding activities stir up bottom sediment and can dramatically decrease water clarity in shallow lakes. As mentioned above, lakes with low water clarity are, generally, considered to have poor water quality. This is because the causes and effects of low clarity negatively impact both plant and fish communities. Fish populations will suffer as water clarity decreases due to a lack of food and decreased ability to successfully hunt for prey. Bluegills are planktivorous fish and feed on invertebrates that inhabit aquatic plants. If low clarity results in the disappearance of plants, this food source will disappear too. Largemouth Bass and Northern Pike are piscivorous fish that feed on other fish and hunt by sight. As the water clarity decreases, these fish species find it more difficult to see and ambush prey and may decline in size as a result. This could eventually lead to an imbalance in the fish community. Phosphorus release from resuspended sediment could increase as water clarity and plant density decrease. This would then result in increased algae blooms, further reducing Secchi depth and aggravating all problems just discussed. The median Secchi depth for Lake County lakes is 2.95 feet. From 2000-2010, both Ozaukee Lake and McDonald Lake #2 had the lowest Secchi depths (0.25 feet) and West Loon Lake had the highest (23.50 feet). As an example of the difference in Secchi depth based on plant coverage, South Churchill Lake, which had no plant coverage and large numbers of Common Carp in 2003 had an average Secchi depth of 0.73 feet (over four times lower than the county average), while Deep Lake, which had a diverse plant community and few carp had an average 2003 Secchi depth of 12.48 feet (almost four times higher than the county average).

Another measure of clarity is the use of a light meter. The light meter measures the amount of light at the surface of the lake and the amount of light at each depth in the water column. The amount of attenuation and absorption (decreases) of light by the water column are major factors controlling temperature and potential photosynthesis. Light intensity at the lake surface varies seasonally and with cloud cover, and decreases with depth. The deeper into the water column light penetrates, the deeper potential plant growth. The maximum depth at which algae and plants can grow underwater is usually at the depth where the amount of light available is reduced

to 0.5%-1% of the amount of light available at the lake surface. This is called the euphotic (sunlit) zone. A general rule of thumb in Lake County is that the 1% light level is about 1 to 3 times the Secchi disk depth.

Alkalinity, Conductivity, Chloride, pH:

Alkalinity:

Alkalinity is the measurement of the amount of acid necessary to neutralize carbonate (CO_3^{2-}) and bicarbonate (HCO_3^-) ions in the water, and represents the buffering capacity of a body of water. The alkalinity of lake water depends on the types of minerals in the surrounding soils and in the bedrock. It also depends on how often the lake water comes in contact with these minerals. If a lake gets groundwater from aquifers containing limestone minerals such as calcium carbonate (CaCO_3) or dolomite (CaMgCO_3), alkalinity will be high. The median alkalinity in Lake County lakes (162 mg/L) is considered moderately hard according to the hardness classification scale of Brown, Skougstad and Fishman (1970). Because hard water (alkaline) lakes often have watersheds with fertile soils that add nutrients to the water, they usually produce more fish and aquatic plants than soft water lakes. Since the majority of Lake County lakes have a high alkalinity they are able to buffer the adverse effects of acid rain.

Conductivity and Chloride:

Conductivity is the inverse measure of the resistance of lake water to an electric flow. This means that the higher the conductivity, the more easily an electric current is able to flow through water. Since electric currents travel along ions in water, the more chemical ions or dissolved salts a body of water contains, the higher the conductivity will be. Accordingly, conductivity has been correlated to total dissolved solids and chloride ions. The amount of dissolved solids or conductivity of a lake is dependent on the lake and watershed geology, the size of the watershed flowing into the lake, the land uses within that watershed, and evaporation and bacterial activity. Many Lake County lakes have elevated conductivity levels in May, but not during any other month. This was because chloride, in the form of road salt, was washing into the lakes with spring rains, increasing conductivity. Most road salt is sodium chloride, calcium chloride, potassium chloride, magnesium chloride or ferrocyanide salts. Beginning in 2004, chloride concentrations are one of the parameters measured during the lake studies. Increased chloride concentrations may have a negative impact on aquatic organisms. Conductivity changes occur seasonally and with depth. For example, in stratified lakes the conductivity normally increases in the hypolimnion as bacterial decomposition converts organic materials to bicarbonate and carbonate ions depending on the pH of the water. These newly created ions increase the conductivity and total dissolved solids. Over the long term, conductivity is a good indicator of potential watershed or lake problems if an increasing trend is noted over a period of years. It is also important to know the conductivity of the water when fishery assessments are conducted, as electroshocking requires a high enough conductivity to properly stun the fish, but not too high as to cause injury or death.

Since 2004 measurements taken in Lake County lakes have exhibited a trend of increasing salinity measured by chloride concentrations. The median near surface chloride concentration of

Lake County Lakes was 142 mg/L. In 2009, Schreiber Lake had the lowest chloride concentration recorded at 2.7 mg/L. The maximum average chloride measurement was at 2760 mg/L at IMC. It is important to note that salt water is denser than fresh water and so it accumulates in the hypolimnion or near the bottom of the lake, this can impact mixing of bottom waters into surface waters in lakes that experience turnover. This phenomenon could have far reaching impacts to an entire ecosystem within a lake. Further, in studies conducted in Minnesota, chloride concentrations as low as 12 mg/L have been found to impact some species of algae.

pH:

pH is the measurement of hydrogen ion (H^+) activity in water. The pH of pure water is neutral at 7 and is considered acidic at levels below 7 and basic at levels above 7. Low pH levels of 4-5 are toxic to most aquatic life, while high pH levels (9-10) are not only toxic to aquatic life they may also result in the release of phosphorus from lake sediment. The presence of high plant densities can increase pH levels through photosynthesis, and lakes dominated by a large amount of plants or algae can experience large fluctuations in pH levels from day to night, depending on the rates of photosynthesis and respiration. Few, if any pH problems exist in Lake County lakes. Typically, the flooded gravel mines in the county are more acidic than the glacial lakes as they have less biological activity, but do not usually drop below pH levels of 7. The median near surface pH value of Lake County lakes was 8.37, with a minimum average of 7.07 in Bittersweet #13 Lake and a maximum of 10.40 in Summerhill Estates Lake.

Eutrophication and Trophic State Index:

The word *eutrophication* comes from a Greek word meaning “well nourished.” This also describes the process in which a lake becomes enriched with nutrients. Over time, this is a lake’s natural aging process, as it slowly fills in with eroded materials from the surrounding watershed and with decaying plants. If no human impacts disturb the watershed or the lake, natural eutrophication can take thousands of years. However, human activities on a lake or in the watershed accelerate this process by resulting in rapid soil erosion and heavy phosphorus inputs. This accelerated aging process on a lake is referred to as cultural eutrophication. The term trophic state refers to the amount of nutrient enrichment within a lake system. *Oligotrophic* lakes are usually deep and clear with low nutrient levels, little plant growth and a limited fishery. *Mesotrophic* lakes are more biologically productive than oligotrophic lakes and have moderate nutrient levels and more plant growth. A lake labeled as *eutrophic* is high in nutrients and can support high plant densities and large fish populations. Water clarity is typically poorer than oligotrophic or mesotrophic lakes and dissolved oxygen problems may be present. A *hypereutrophic* lake has excessive nutrients, resulting in nuisance plant or algae growth. These lakes are often pea-soup green, with poor water clarity. Low dissolved oxygen may also be a problem, with fish kills occurring in shallow, hypereutrophic lakes more often than less enriched lakes. As a result, rough fish (tolerant of low dissolved oxygen levels) dominate the fish community of many hypereutrophic lakes. The categorization of a lake into a certain trophic state should not be viewed as a “good to bad” categorization, as most lake residents rate their

lake based on desired usage. For example, a fisherman would consider a plant-dominated, clear lake to be desirable, while a water-skier might prefer a turbid lake devoid of plants. Most lakes in Lake County are eutrophic or hypereutrophic. This is primarily as a result of cultural eutrophication. However, due to the fertile soil in this area, many lakes (especially man-made) may have started out under eutrophic conditions and will never attain even mesotrophic conditions, regardless of any amount of money put into the management options. This is not an excuse to allow a lake to continue to deteriorate, but may serve as a reality check for lake owners attempting to create unrealistic conditions in their lakes.

The Trophic State Index (TSI) is an index which attaches a score to a lake based on its average total phosphorus concentration, its average Secchi depth (water transparency) and/or its average chlorophyll *a* concentration (which represent algae biomass). It is based on the principle that as phosphorus levels increase, chlorophyll *a* concentrations increase and Secchi depth decreases. The higher the TSI score, the more nutrient-rich a lake is, and once a score is obtained, the lake can then be designated as oligotrophic, mesotrophic or eutrophic. Table 1 (below) illustrates the Trophic State Index using phosphorus concentration and Secchi depth.

Table 1. Trophic State Index (TSI).

Trophic State	TSI score	Total Phosphorus (mg/L)	Secchi Depth (feet)
Oligotrophic	<40	≤ 0.012	>13.12
Mesotrophic	≥40<50	>0.012 ≤ 0.024	≥6.56<13.12
Eutrophic	≥50<70	>0.024 ≤ 0.096	≥1.64<6.56
Hypereutrophic	≥70	>0.096	< 1.64